

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problems Mailbox.**

THIS PAGE BLANK (USPTO)



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 14/31, 16/12, C12N 15/31, G01N 33/577, A61K 39/085, 39/09		A1	(11) International Publication Number: WO 98/45325
			(43) International Publication Date: 15 October 1998 (15.10.98)
(21) International Application Number: PCT/US98/06663		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 1 April 1998 (01.04.98)			
(30) Priority Data: 08/838,413 7 April 1997 (07.04.97) US			
(71) Applicant: THE ROCKEFELLER UNIVERSITY [US/US]; 1230 York Avenue, New York, NY 10021-6399 (US).			
(72) Inventors: BANNAN, Jason, D.; 2600 Milton Lane, Thompson Station, TN 37179 (US). ZABRISKIE, John, B.; 1385 York Avenue, New York, NY 10021 (US).			
(74) Agents: SMITH, Stephen, R. et al.; Morgan & Finnegan, L.L.P., 345 Park Avenue, New York, NY 10154 (US).		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	

(54) Title: PEPTIDES USEFUL FOR REDUCING SYMPTOMS OF TOXIC SHOCK SYNDROME

TOXIN	REGION 1	REGION 2
PEPs :	CMYGGVTEHEGN	KKNVTVOELDYKIRKYLVDNKKLY
SEA	130 CMYGGVTLHDNN 141	171 KKNVTVOELDLQARRYLQEKYNLY 194
SEB	140 CMYGGVTEHNGN 151	179 KKKVTAQELDYLTRHYLVKNKKLY 202
SEC	137 CMYGGITKHEGN 148	178 KKSMTAQELDIKARNFLINKKNLY 201
SED	131 CTYGGVTPHEGN 142	172 KKNVTVOELDAQARRYLQKDLKLY 195
SEE	130 CMYGGVTLHDNN 141	171 KKEVTVOELDIQARHYLHGKFGLY 194
SEH	116 CLYGGITL.NSE 126	151 KKNVTLQELDIKIRKILSDKYKIY 174
SPEA	128 CIYGGVTNHEGN 139	167 KKMVTAQELDYKVRKYLTDNKQLY 190
SPEC	112 YIYGGITPAQNN 123	151 KDIVTFQELDFKIRKLYMDNYKIY 174
SSA	134 CMYGGVTEHHRN 145	174 KKQVTVOELDCRTRKILVSRKNLY 197
TSST1		161 KKQLAISTLDFEIRHQLTQIHGLY 184

(57) Abstract

This invention relates to compositions and methods for eliciting an immunogenic response in mammals, including responses which provide protection against, or reduce the severity, of toxic shock from bacterial infections. More particularly it relates to peptides derived from homologous sequences of the family of staphylococcal and streptococcal toxins, which may be polymeric, and carrier-conjugates thereof, and their use to induce serum antibodies. The invention also relates to serum antibodies induced by the peptides and carrier-conjugates and their use to prevent, treat, or protect against the toxic effects of most, if not all, of the staphylococcal and streptococcal toxins. The invention also relates to diagnostic assays and kits to detect the presence of staphylococcal and streptococcal toxins, or antibodies thereto. The invention also relates to isolated and purified nucleic acids encoding the peptides of the invention and transformed host cells containing those nucleic acids.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

-1-

PEPTIDES USEFUL FOR REDUCING SYMPTOMS
OF TOXIC SHOCK SYNDROME

FIELD OF THE INVENTION

This invention relates to compositions and methods for eliciting an immunogenic response in mammals, including responses which provide protection against, or reduce the severity, of toxic shock syndrome from bacterial infections. More particularly it relates to peptides, which may be polymeric, and carrier-conjugates thereof, derived from homologous sequences of the family of staphylococcal and streptococcal pyrogenic toxins. The peptides of the invention are useful to induce serum antibodies and may also be useful in diagnostic assays.

The invention also relates to antibodies induced by the peptides and/or carrier-conjugates and their use to prevent, treat, or protect against the toxic effects of bacterial toxins, including most, if not all, of the staphylococcal and streptococcal pyrogenic toxins. The invention also relates to compositions and methods to protect against, or ameliorate the effects of, autoimmune diseases which are associated with, or are the result of, the presence of staphylococcal or streptococcal toxins.

The invention also relates to diagnostic assays and kits to detect the presence of staphylococcal and streptococcal pyrogenic toxins, or antibodies thereto.

The invention also relates to isolated and purified nucleic acids encoding the peptides of the invention and transformed host cells containing those nucleic acids.

BACKGROUND OF THE INVENTION

The pyrogenic exotoxins of Group A streptococci and the enterotoxins of Staphylococcus aureus, which are also pyrogenic exotoxins, constitute a family of structurally related toxins which share similar biological activities (11, 13). The staphylococcal and streptococcal pyrogenic

-2-

exotoxins also share significant amino acid homology throughout their sequences (11, 19, 40). This pyrogenic exotoxin family contains nine main toxin types, and several allelic variants (subtypes) have been described. Several studies have shown that the toxins share common motifs based on immunologic cross reactivity between the toxins (26, 27). These toxins share the ability to bind the major histocompatibility complex (MHC) molecules of infected hosts, as well as the variable beta chain of the T-cell receptor complex (TCR), causing an aberrant proliferation of specific T-cell subsets (3, 4, 12). This property of the toxins has labeled them as "superantigens" (36) since they do not interact with the MHC and TCR molecules in the manner of conventional antigens (14, 18).

These bacterial toxins cause a variety of syndromes in humans. Staphylococcal enterotoxins have been implicated in staphylococcal food poisoning (26), as well as toxic shock like syndromes (1). The gene sequences and deduced amino acid sequences of at least six staphylococcal enterotoxins ("SE"): A, B, C, D, E and H, are known, i.e., SEA, SEB, SEC, SED, SEE, and SEH (19, 23). The streptococcal pyrogenic exotoxins ("SPE") have been implicated in causing the symptoms of scarlet fever and toxic shock like syndrome (8, 20, 30). The sequences of three members of this family are known: SPEA, SPEC, and SSA (5, 23, 35).

Toxic shock syndrome toxin (TSST-1) from S. aureus shares similar biological activity with the enterotoxins and streptococcal pyrogenic exotoxins, however it is not as closely related structurally (2). Toxic shock syndrome can be exacerbated by the synergistic effects of TSST-1 with the enterotoxin/pyrogenic toxin family of toxins (9, 25). Gram negative bacterial endotoxin and the pyrogenic toxins can work synergistically to produce lethal toxic shock (17, 30).

"Toxic shock like syndrome" is the term previously

-3-

used to describe the syndromes caused by staphylococcal and streptococcal pyrogenic bacterial exotoxins other than toxic shock syndrome toxin (TSST-1) from S. aureus. Currently, the term "toxic shock syndrome" is used to describe the syndromes caused by TSST-1 and the other pyrogenic exotoxins, and is the terminology used hereinafter.

SUMMARY OF THE INVENTION

The present invention relates to the identification of consensus sequences derived from two conserved regions of the staphylococcal enterotoxins and streptococcal pyrogenic toxins (hereinafter called "region 1" and "region 2") and the discovery that compositions comprising amino acid sequences based on these two conserved regions of the staphylococcal enterotoxins and streptococcal pyrogenic exotoxins are capable of inducing antibodies which react with a variety of staphylococcal and streptococcal pyrogenic exotoxins and are also capable of ameliorating or preventing diseases related to the deleterious effects of these toxins.

The invention also relates to compositions and methods for preventing and treating diseases related to the release of certain pyrogenic exotoxins from bacteria.

This invention provides amino acid sequences capable of inducing antibodies that reduce, inhibit or eliminate the deleterious effects of bacterial toxins, such as those of staphylococcus and a variety of streptococci. These antibodies may be induced by administration of a pharmaceutical composition and/or vaccine containing a composition comprising a peptide derived from one or both of the two conserved regions described herein, or a structurally and/or immunologically related antigen.

The amino acid sequences provided by this invention are sufficiently common to all members of this family of pyrogenic exotoxins to be useful for eliciting antibodies

-4-

- o which are cross-reactive with toxins derived from various bacteria.

5 The amino acid sequences provided by this invention are also useful for new methods of preventing and treating symptoms associated with the bacterial release of the staphylococcal enterotoxins and the streptococcal pyrogenic exotoxins. Such methods include, for example, administering to an individual at risk of infection or developing a toxic reaction to the exotoxins at least one of the consensus amino acid sequences of this invention in an amount sufficient to elicit the production of antibodies to the exotoxins.

10 In a preferred embodiment of this invention, an individual at risk for developing toxic shock syndrome or an individual with symptoms of toxic shock syndrome may be treated by administering to such individual antibodies which have been generated in a mammal immunized with at least one of the compositions of this invention.

15 Vaccines and pharmaceutical compositions comprising at least one of the consensus amino acid sequences and a physiologically acceptable carrier and optionally an adjuvant are also part of this invention.

20 Another object of the invention is to provide antibodies induced by the peptides and carrier-conjugates thereof. These antibodies may be used to prevent, treat, or protect against the toxic effects of most, if not all, of the staphylococcal and streptococcal pyrogenic exotoxins. The antibodies may also be useful to protect against, or ameliorate the effects of, autoimmune diseases which are associated with, or are the result of, the presence of staphylococcal or streptococcal pyrogenic exotoxins. These antibodies are also useful in diagnostic assays and kits to detect the presence of staphylococcal and streptococcal pyrogenic exotoxins and to aid in the diagnosis of diseases related to the presence of those toxins.

35

-5-

Another object of the invention is to provide isolated and purified nucleic acids encoding the amino acid sequences of the invention, as well as suitable expression systems, vector components and transformed host cells containing those nucleic acids.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. Comparison of the synthetic peptide sequences to conserved regions 1 and 2 of the staphylococcal enterotoxins (SEA, SEB, SEC, SED, SEE, and SEH), and streptococcal pyrogenic exotoxins (SPEA, SPEC, and SSA). Staphylococcal toxic shock syndrome toxin 1 (TSST-1) was compared with the region 2 peptide. Numbers represent the residue positions as a reference to where these regions exist in the whole toxin molecules. Sequences are from either the Swiss protein or Genbank databases under the following accession numbers. Swiss protein: SPEA, P08095; SPEC, P13380; SEA, P13163; SEB, P01552; SEC, P01553; SED, P20723; SEE, P12993. Genbank: SEH, U11702; SSA, L29565; TSST1, J02615.

Figure 2. ELISA titers of antibodies from rabbits immunized with polymeric peptide #6348. The peptide was diluted so that it was delivered to each well to give a final concentration of 2 $\mu\text{g}/100 \mu\text{l}$. The serum was then diluted to 1:1,000; 1:10,000; 1:100,000; 1:500,000; and 1:1,000,000 and 100 μl of each dilution of serum was placed in each well. Experiments were run in triplicate for each dilution of serum. Note the 1 log higher titers of rabbit #443 serum as compared to rabbit #442 serum. Cut off readings were at O.D. 0.6.

Figure 3. 12% SDS PAGE gels and immunoblot of a variety of staphylococcal and streptococcal toxins developed with the anti-peptide 6348 antibody. Note bands of correct molecular weight (M.W.) of each toxin identified by the anti-peptide antibody. Lane 1: SPEA, lane 2: SEA, lane 3: SEB, lane 4: SED, lane 5: SEE, lane

-6-

- 6: SEC and lane 7 TsST-1. Note bands at appropriate M.W. in lanes 1-4. Fainter bands are seen in lanes 5 and 7.

Figure 4. Bar graphs of blastogenesis assays of human mononuclear cell populations stimulated by various toxins in the presence of normal rabbit serum and anti-peptide 6348 serum. Note the marked inhibition of SEB, SEC, SEE, SPEA and SPEC by the anti-peptide antibody. Less, but definite, inhibition of SEA by the anti-peptide antibody was also seen.

DETAILED DESCRIPTION OF THE INVENTION

It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only, and are not restrictive of the invention, as claimed. The accompanying drawings, which are incorporated in and constitute a part of the specification, illustrate an embodiment of the invention and, together with the description, serve to explain the principles of the invention.

Two consensus patterns, corresponding to conserved region 1 and region 2, respectively, are identified as common to members of the staphylococcal enterotoxin and streptococcal pyrogenic toxin family of toxins when the program "Motifs" in a software package from the Genetics Computer Group, Inc. ("GCG") is run using the streptococcal SPEC toxin as an example. "Program Manual for the Wisconsin Package, Version 8, September 1994, Genetics Computer Group, 575 Science Drive, Madison, Wisconsin, USA 53711", incorporated herein by reference.

The first consensus sequence ("GCG consensus #1") identified by the Motifs program has the amino acid sequence YGG(LIV)TXXXXN, which is rewritten herein as YGGX₁TX₂X₃X₄X₅N (SEQ ID NO:1), wherein X₁ is selected from the group consisting of L, I, or V; and X₂, X₃, X₄ and X₅ are each independently selected from the group consisting

-7-

of any amino acid. This pattern is present in the staphylococcal enterotoxins and streptococcal pyrogenic exotoxins, but not in TSST-1. The second consensus sequence ("GCG consensus #2") identified by the Motifs program has the amino acid sequence

KXX(LIV)XXXX(LIV)DXXXRXXLXXXXX(LIV)Y, rewritten herein as
KX₆X₇X₈X₉X₁₀X₁₁X₁₂X₁₃DX₁₄X₁₅X₁₆RX₁₇X₁₈LX₁₉X₂₀X₂₁X₂₂X₂₃X₂₄Y (SEQ ID NO: 2), wherein X₈, X₁₃ and X₂₄ are each independently selected from the group consisting of L, I and V, and X₆, X₇, X₉, X₁₀, X₁₁, X₁₂, X₁₄, X₁₅, X₁₆, X₁₇, X₁₈, X₁₉, X₂₀, X₂₁, X₂₂ and X₂₃ are each independently selected from the group consisting of any amino acid. This pattern is present in the staphylococcal enterotoxins, streptococcal pyrogenic exotoxins, and TSST-1.

One object of the invention is to provide compositions comprising peptides comprising amino acid sequences based on these two conserved regions of the staphylococcal enterotoxins and streptococcal pyrogenic toxins. These peptides may be used for eliciting an immunogenic response in mammals, including responses which provide protection against, or reduce the severity, of toxic shock from staphylococcal or streptococcal infections. These peptides may also be useful to protect against, or ameliorate the effects of, autoimmune diseases which are associated with, or are the result of, the presence of staphylococcal or streptococcal pyrogenic exotoxins. These peptides are also useful in diagnostic assays and kits to detect the presence of antibodies to staphylococcal and streptococcal pyrogenic exotoxins and to aid in the diagnosis of diseases related to the presence of those toxins.

The peptides of the invention are those derived from either one or both of the following two consensus sequences:

YGGX₁TX₂X₃X₄X₅N (SEQ ID NO:1), wherein X₁ is selected from the group consisting of L, I, or V; and X₂, X₃, X₄ and X₅

-8-

are each independently selected from the group consisting of any amino acid.

KXX(LIV)XXXX(LIV)DXXXRXXLXXXXX(LIV)Y, rewritten herein as $KX_6X_7X_8X_9X_{10}X_{11}X_{12}X_{13}DX_{14}X_{15}X_{16}RX_{17}X_{18}LX_{19}X_{20}X_{21}X_{22}X_{23}X_{24}Y$ (SEQ ID NO: 2), wherein X_8 , X_{13} and X_{24} are each independently selected from the group consisting of L, I and V, and X_6 , X_7 , X_9 , X_{10} , X_{11} , X_{12} , X_{14} , X_{15} , X_{16} , X_{17} , X_{18} , X_{19} , X_{20} , X_{21} , X_{22} and X_{23} are each independently selected from the group consisting of any amino acid.

A preferred consensus sequence of the invention from region 1 (consensus #1a) has the amino acid sequence $X_{25}X_{26}YGGX_1TX_2X_3X_4X_5N$ (SEQ ID NO: 28), wherein X_1 is selected from the group consisting of L, I, and V; X_2 , X_4 and X_5 are each independently selected from the group consisting of any amino acid; and X_3 , X_{25} and X_{26} are each independently selected from the group consisting of any amino acid and of no amino acid; but preferably X_1 is selected from the group consisting of I and V; X_2 is selected from the group consisting of L, E, K, P and N; X_3 is selected from the group consisting of H and A and no amino acid; X_4 is selected from the group consisting of D, N, E, Q, and H; X_5 is selected from the group consisting of N, G, S, and R; X_{25} is selected from the group consisting of C and Y and no amino acid; and X_{26} is selected from the group consisting of M, T, L, I, and no amino acid.

A preferred consensus sequence of the invention from region 2 (consensus #2a) has the amino acid sequence: $KX_6X_7X_8X_9X_{10}X_{11}X_{12}X_{13}DX_{14}X_{15}X_{16}RX_{17}X_{18}X_{27}X_{19}X_{20}X_{21}X_{22}X_{23}X_{24}Y$ (SEQ ID NO: 29), wherein X_8 , X_{13} and X_{24} are each independently selected from the group consisting of L, I and V; X_6 , X_7 , X_9 , X_{10} , X_{11} , X_{12} , X_{14} , X_{15} , X_{16} , X_{17} , X_{18} , X_{19} , X_{20} , X_{21} , X_{22} , and X_{23} are each independently selected from the group consisting of any amino acid; and X_{27} is selected from the group consisting of L and Y; but preferably X_6 is selected from the group consisting of K and D; X_7 is selected from the group consisting of N, K, S, E, M, I and Q; X_8 is selected

35

-9-

from the group consisting of L and V; X₉ is selected from the group consisting of T and A; X₁₀ is selected from the group consisting of V, A, L, F and I; X₁₁ is selected from the group consisting of Q and S; X₁₂ is selected from the group consisting of E and T; X₁₃ is selected from group consisting of L and I; X₁₄ is selected from the group consisting of L, Y, I, A, F and C; X₁₅ is selected from the group consisting of Q, L, K and E; X₁₆ is selected from the group consisting of A, T, I and V; X₁₇ is selected from the group consisting of R, H, N and K; X₁₈ is selected from the group consisting of Y, F, I, L and Q; X₁₉ is selected from the group consisting of Q, V, I, H, S, T and M; X₂₀ is selected from the group consisting of E, K, N, G, D, S and Q; X₂₁ is selected from the group consisting of K, N, D, R and I; X₂₂ is selected from the group consisting of Y, K, L, F and H; X₂₃ is selected from the group consisting of N, K, G and Q; X₂₄ is selected from the group consisting of L and I; and X₂₇ is L.

The following Table 1 lists the amino acids that are found at each of the variable positions in the sequences shown in Figure 1, and the number of times they appear at that position:

25

30

35

-10-

Table 1

Frequency of the amino acids in the variable positions in
the sequences shown in Figure 1

	X ₁	6V	3I					
5	X ₂	3L	2E	1K	2P	1N		
	X ₃	7H	1A	one deletion (no amino acid)				
	X ₄	2D	2N	3E	1Q	1H		
	X ₅	3N	4G	1S	1R			
	X ₆	9K	1D					
10	X ₇	3N	1K	1S	1E	1M	1I	1Q
	X ₈	9V	1L					
	X ₉	9T	1A					
	X ₁₀	4V	3A	1L	1F	1I		
	X ₁₁	9Q	1S					
15	X ₁₂	9E	1T					
	X ₁₃	9L	1I					
	X ₁₄	2L	2Y	2I	1A	2F	1C	
	X ₁₅	3Q	1L	5K	1E			
20	X ₁₆	4A	2T	3I	1V			
	X ₁₇	2R	3H	1N	4K			
	X ₁₈	5Y	1F	2I	1L	1Q		
	X ₁₉	2Q	2V	1I	1H	1S	2T	1M
	X ₂₀	1E	2K	1N	1G	3D	1S	1Q
25	X ₂₁	4K	3N	1D	1R	1I		
	X ₂₂	3Y	4K	1L	1F	1H		
	X ₂₃	3N	4K	2G	1Q			
	X ₂₄	8L	2I					
	X ₂₅	8C	1Y					
30	X ₂₆	5M	2I	1L	1T			
	X ₂₇	9L	1Y					

In the peptides of the present invention, X₁, X₈, X₁₃
and X₂₄ may each independently be selected from the group
consisting of L, I and V; X₂, X₃, X₄, X₅, X₆, X₇, X₉, X₁₀,

-11-

0 X₁₁, X₁₂, X₁₄, X₁₅, X₁₆, X₁₇, X₁₈, X₁₉, X₂₀, X₂₁, X₂₂, X₂₃, X₂₅ and
X₂₆ may each independently be any amino acid; X₃, X₂₅ and
X₂₆ may also each independently be no amino acid; and X₂₇
is selected from the group consisting of L and Y.
However, in general, the amino acids present at the
positions X₁ to X₂₇ in the toxins shown in Figure 1 (and
5 listed in Table 1) are preferred for those positions, and
the amino acids present most often at those positions in
the toxins shown in Figure 1 (and listed in Table 1) are
more preferred. For example, from Figure 1, and Table 1,
10 it can be determined that H (histidine) is present in
seven toxins at position X₃ and A (alanine) is present in
one toxin at position X₃, and there is no amino acid
present in one toxin at X₃. These are the preferred amino
acids for position X₃. The more preferred amino acid for
15 position X₃ in a peptide of the invention is H
(histidine). The more preferred amino acids for X₁
through X₂₆ are: X₁ = valine; X₂ = leucine; X₃ = histidine;
X₄ = glutamic acid; X₅ = glycine; X₆ = lysine; X₇ =
asparagine; X₈ = valine; X₉ = threonine; X₁₀ = valine; X₁₁ =
glutamine; X₁₂ = glutamic acid; X₁₃ = leucine; X₁₄ =
20 leucine, tyrosine, isoleucine or phenylalanine; X₁₅ =
lysine; X₁₆ = alanine; X₁₇ = lysine; X₁₈ = tyrosine; X₁₉ =
glutamine, valine or threonine; X₂₀ = aspartic acid; X₂₁ =
lysine; X₂₂ = lysine; X₂₃ = lysine; X₂₄ = leucine; X₂₅ =
cysteine; X₂₆ = methionine; and X₂₇ = leucine. But note
25 that in the exemplified peptides of the invention
described hereinbelow, i.e., SEQ ID NOS: 6, 7 and 8,
inosine (I) is used at position X₁₆ instead of the more
frequently found alanine (A).

30 As is evident from Figure 1 and the above Table 1,
some amino acid residues are much more highly conserved
than suggested by the GCG package data provided by the
"Motifs" program.

35 In region 1, the preferred consensus is larger
(consensus #1a), and usually includes a C in the first

-12-

- position (X_{25}). The second residue (X_{26}) is most often a M, but this can vary. In the ninth position (X_3), H is the most highly conserved. The eleventh residue (X_5) is most often a G.

In region 2, the preferred consensus (consensus #2a) is much more highly conserved than suggested by the GCG program, especially if one excludes TSST-1 sequences from consideration, as follows: The second position (X_6) is more highly conserved than suggested, being almost exclusively a K; the fourth residue (X_8) is always a V followed exclusively by a T in the fifth position (X_9); the sixth position (X_{10}) is somewhat variable; but the seventh position (X_{11}) is always a Q, followed by E (X_{12}). The next position is almost always an L (X_{13}), and the second to last position (X_{24}) is almost always an L.

Thus, additional modified consensus sequences for region 1 and region 2, which are of narrower scope than the GCG consensus sequences #1 and #2 and the modified consensus sequences #1a and #2a, are as follows:

Consensus #1b:

CMYGGX₁TX₂HX₄GN (SEQ ID NO: 30)

wherein

X_1 is V or I, preferably V;

X_2 is L, E, K, P or N, preferably E or L;

and

X_4 is D, N, E, Q or H, preferably E.

Consensus #2b:

KKX₇VTX₁₀QELDX₁₄X₁₅X₁₆RX₁₇X₁₈X₂₇X₁₉X₂₀X₂₁X₂₂X₂₃LY (SEQ ID NO:31)

wherein

X_7 is N, K, S, E, M, I or Q, preferably N;

X_{10} is V, A, L, F or I, preferably V;

X_{14} is L, Y, I, A, F or C, preferably Y;

X_{15} is Q, L, K or E, preferably K;

X_{16} is A, T, I or V, preferably I;

X_{17} is R, H, N or K, preferably K;

-13-

o X₁₈ is Y, F, I, L or Q, preferably Y;
X₁₉ is Q, V, I, H, S, T or M, preferably V;
X₂₀ is E, K, N, D, G, S or Q, preferably D;
X₂₁ is K, N, D, R or I, preferably N;
X₂₂ is Y, K, L, F or H, preferably K;
5 X₂₃ is N, K, G or Q, preferably K; and
X₂₇ is L or Y, preferably L.

Peptides exemplified herein are CMYGGVTEHEGN (SEQ ID NO: 3), CMYGGVTEHEGNGC* (SEQ ID NO: 5), KKNVTVQELDYKIRKYLVDNKKLY (SEQ ID NO: 4), CGKKNVTVQELDYKIRKYLVDNKKLYGC* (SEQ ID NO: 6),
10 CMYGGVTEHEGNKKNVTVQELDYKIRKYLVDNKKLY (SEQ ID NO: 7) and CMYGGVTEHEGNKKNVTVQELDYKIRKYLVDNKKLYGC* (SEQ ID NO: 8), wherein an asterisk indicates that the peptide is a randomly cross-linked polymer. The exemplified polymer peptides are at least 6,000 to 8,000 daltons. The average
15 size of the exemplified polymer peptides is about 12,000 to 15,000 daltons. Small peptides and/or contaminants may be removed by dialysis or other methods available in the art. Similarly, larger aggregates may be removed using, e.g., a 0.25 micron filter, which can also be used to
20 sterilize the peptides.

Note that the amino acids cysteine and methionine, "CM", are present at the amino terminus of the exemplified region 1 peptides since those amino acids are most often found in that position in nature. Note also that the
25 amino acids cysteine and glycine, "CG" and "GC", are used at the amino and/or carboxy- termini of some of the exemplified region 2 peptides. The amino acid cysteine "C" is used to facilitate cross-linking through the formation of disulfide bonds. The amino acid glycine,
30 "G", is used as a spacer residue.

The preferred peptides of the invention are those which exclude full length native toxin molecules. The preferred peptides of this invention are not toxic, but toxic peptides maybe useful in this invention, for

35

-14-

example, in eliciting antibodies in a non-human system. The most preferred peptides of the invention do not contain amino acid sequences in the sequence in which they are found in any particular native toxin molecule.

The present invention encompasses monomers of the peptides derived from either one or both of the two consensus regions described herein. These monomers may comprise one or more sequences derived from either region 1 or region 2 or both, such as consensus sequences #1 and #2, preferably consensus sequences #1a and/or #2a, more preferably consensus sequences #1b and/or #2b, most preferably one or more of the exemplified consensus sequence peptides. If the monomer contains more than one consensus sequence, these sequences may be immediately adjacent to each other or separated by a linker. In addition, different orientations of the peptides are within the scope of this invention. Furthermore, the order of the consensus peptides within the full peptide may be variable.

The present invention also encompasses homogeneous or heterogeneous polymers of the peptides disclosed herein (e.g., concatenated, cross-linked and/or fused identical peptide units or concatenated, cross-linked and/or fused diverse peptide units), and mixtures of the peptides, polymers, and/or conjugates thereof.

Linkers useful in the invention may, for example, be simply peptide bonds, or may comprise amino acids, including amino acids capable of forming disulfide bonds, but may also comprise other molecules such as, for example, polysaccharides or fragments thereof.

In the peptides exemplified herein, sequences derived from consensus region 1 and consensus region 2 may be immediately adjacent to each other, linked by peptide bonds, (see, e.g., SEQ ID NO:7) and/or connected via amino acid linkers capable of forming di-sulfide bonds via cysteine residues (see, e.g., SEQ ID NO: 8). In the

-15-

native toxin molecules, the sequences of region 1 and region 2 are separated by 27 amino acids. When the linkers are additional amino acids, they are most preferably 1 to 27 amino acids in length, although longer linkers may also be used in accordance with this invention.

The linkers for use with this invention may be chosen so as to contribute their own immunogenic effect which may be either the same, or different, than that elicited by the consensus sequences of the invention. For example, such linkers may be bacterial antigens which also elicit the production of antibodies to infectious bacteria. In such instances, for example, the linker may be a protein or protein fragment of an infectious bacteria, or a bacterial polysaccharide or polysaccharide fragment.

A peptide of the invention includes any substituted analog or chemical derivative of a peptide derived from one or both of the two consensus regions described herein, most preferably of the exemplified peptides described herein, so long as the peptide is capable of either eliciting the production of antibodies capable of binding to most of the staphylococcal and streptococcal pyrogenic exotoxins, or reacting with (i.e., specifically binding to) antibodies that react with most of the staphylococcal and streptococcal pyrogenic exotoxins. Therefore, a peptide can be subject to various changes that provide for certain advantages in its use.

The peptides of the invention are useful for providing active immunization for the prevention of disease related to the deleterious effects of staphylococcal and streptococcal pyrogenic exotoxins and for preparation of antibodies as a passive immunization therapy. The peptides are designed to induce antibodies which react with a variety of staphylococcal and streptococcal pyrogenic exotoxins (preferably with at least two, more preferably with at least four, and most

35

-16-

preferably with at least seven of the pyrogenic exotoxins) for use in therapy to increase resistance to, prevent and/or treat toxic shock syndrome. The peptides may also be useful to protect against, or ameliorate the effects of, autoimmune diseases which are associated with, or are the result of, the presence of staphylococcal or streptococcal exotoxins. The peptides of the invention will also be useful in diagnostic tests for detecting antibodies to staphylococcal and streptococcal pyrogenic exotoxins.

The peptide may be mixed with an adjuvant. The peptide also may be bound to a non-toxic non-host protein carrier to form a conjugate or it may be bound to a saccharide carrier and/or a non-toxic non-host protein carrier to form a conjugate.

The molecular weight of the peptide monomers having one consensus sequence of the invention range from about 1000 to 5000 daltons. Such lower molecular weight species of the invention may be useful as immunogens themselves or, more preferably, may be used as haptens conjugated to a larger carrier molecule, such as, for example, a protein. As with other peptides, the molecular weight of the peptide alone, when conjugated to a carrier or in the presence of an adjuvant, is related to its immunogenicity. Thus, the peptide may vary in molecular weight in order to enhance its antigenicity or immunogenicity. In an exemplified embodiment, the molecular weight of the peptide, in polymeric form, is greater than about 6000 to 8000 daltons, with an average weight of 12,000 to 15,000 daltons. The total size of the peptide is only limited to its ability to be physiologically tolerated.

The invention also relates to isolated and purified nucleic acid molecules which code for the peptides of the invention. The encoded peptides may be monomers, polymers or linked to other peptide sequences (i.e., they may be

-17-

fusion proteins). Other features of the invention include vectors which comprise the nucleic acid molecules of the invention operably linked to promoters, as well as cell lines, such as prokaryotic (e.g., E. coli) and eukaryotic (e.g., CHO and COS) cells transfected with the nucleic acid molecules of the invention. Vectors and compositions for enabling production of the peptides in vivo, i.e., in the individual to be treated or immunized, are also within the scope of this invention.

The nucleic acids encoding the peptides of the invention can be introduced into a vector such as a plasmid, cosmid, phage, virus or mini-chromosome and inserted into a host cell or organism by methods well known in the art. In general, the vectors containing these nucleic acids can be utilized in any cell, either eukaryotic or prokaryotic, including mammalian cells (e.g., human (e.g., HeLa), monkey (e.g., Cos), rabbit (e.g., rabbit reticulocytes), rat, hamster (e.g., CHO and baby hamster kidney cells) or mouse cells (e.g., L cells), plant cells, yeast cells, insect cells or bacterial cells (e.g., E. coli). The vectors which can be utilized to clone and/or express these nucleic acids are the vectors which are capable of replicating and/or expressing the nucleic acids in the host cell in which the nucleic acids are desired to be replicated and/or expressed. See, e.g., F. Ausubel et al., Current Protocols in Molecular Biology, Greene Publishing Associates and Wiley-Interscience (1992) and Sambrook et al. (1989) for examples of appropriate vectors for various types of host cells. Strong promoters compatible with the host into which the gene is inserted may be used. These promoters may be inducible. The host cells containing these nucleic acids can be used to express large amounts of the protein useful in pharmaceuticals, diagnostic reagents, vaccines and therapeutics.

The nucleic acids could be used, for example, in the

-18-

production of peptides for diagnostic reagents, vaccines and therapies for pyrogenic exotoxin related diseases. For example, vectors expressing high levels of peptide can be used in immunotherapy and immunoprophylaxis, after expression in humans. Such vectors include retroviral vectors and also include direct injection of DNA into muscle cells or other receptive cells, resulting in the efficient expression of the peptide, using the technology described, for example, in Wolff et al., Science 247:1465-1468 (1990), Wolff et al., Human Molecular Genetics 1(6):363-369 (1992) and Ulmer et al., Science 259:1745-1749 (1993).

In another embodiment of this invention antibodies are provided which react with peptides of the invention, as well as a variety of staphylococcal and streptococcal pyrogenic exotoxins (preferably with at least two, more preferably with at least four, and most preferably with at least seven of the pyrogenic exotoxins). These antibodies will be useful for passive immunization therapy to increase resistance to or prevent toxic shock syndrome or other diseases related to the presence of bacterial pyrogenic exotoxin. The antibodies may also be useful to protect against, or ameliorate the effects of, autoimmune diseases which are associated with, or are the result of, the presence of staphylococcal or streptococcal pyrogenic exotoxins. The antibodies of the invention will also be useful in diagnostic tests and kits for detecting the presence of staphylococcal and streptococcal pyrogenic exotoxins. These uses are discussed in more detail below.

Methods for preparing peptides of the invention

The peptides of the invention may be prepared by synthetic methods or by recombinant DNA methods, as known in the art and as described herein.

Pharmaceutical Compositions

The pharmaceutical compositions of this invention contain an effective, immunogenic amount of peptide of

-19-

0 this invention. The effective amount of peptide per unit
dose sufficient to induce an immune response depends,
among other things, on the species of mammal inoculated,
the body weight of the mammal and the chosen inoculation
regimen, as well as the presence or absence of an
5 adjuvant, as is well known in the art. Inocula
typically contain peptide concentrations of about 1
microgram to about 1000 micrograms per inoculation (dose),
preferably about 3 micrograms to about 100 micrograms per
dose, most preferably about 5 micrograms to 50 micrograms.

10 The term "unit dose" as it pertains to the inocula
refers to physically discrete units suitable as unitary
dosages for mammals, each unit containing a predetermined
quantity of active material (peptide) calculated to
produce the desired immunogenic effect in association with
the required diluent.

15 Inocula are typically prepared as a solution in a
physiologically acceptable carrier such as saline,
phosphate-buffered saline and the like to form an aqueous
pharmaceutical composition.

20 The route of inoculation of the peptides of the
invention is typically parenteral and is preferably
intramuscular, sub-cutaneous and the like, which results
in eliciting antibodies protective against the deleterious
effects of staphylococcal and streptococcal pyrogenic
exotoxins. The dose is administered at least once. In
25 order to increase the antibody level, at least one booster
dose may be administered after the initial injection,
preferably at about 4 to 6 weeks after the first dose.
Subsequent doses may be administered as indicated.

30 To monitor the antibody response of individuals
administered the compositions of the invention, antibody
titers may be determined. In most instances it will be
sufficient to assess the antibody titer in serum or plasma
obtained from such an individual. Decisions as to whether
to administer booster inoculations or to change the amount

35

-20-

o of the composition administered to the individual may be at least partially based on the titer.

5 The titer may be based on either an immunobinding assay which measures the concentration of antibodies in the serum which bind to a specific antigen, i.e. peptide or toxin; or bactericidal assays which measure the ability of the antibodies to participate with complement in killing bacteria. The ability to neutralize in vitro and in vivo biological effects of the pyrogenic exotoxins may also be assessed to determine the effectiveness of the treatment. See Example 1.

10 Antibodies

The term "antibodies" is used herein to refer to immunoglobulin molecules and immunologically active portions of immunoglobulin molecules. Exemplary antibody molecules are intact immunoglobulin molecules, substantially intact immunoglobulin molecules and portions of an immunoglobulin molecule, including those portions known in the art as Fab, Fab', F(ab')₂ and F(v) as well as chimeric antibody molecules.

20 An antibody of the present invention is typically produced by immunizing a mammal with an immunogen or vaccine containing one or more peptides of the invention, or a structurally and/or antigenically related molecule, to induce, in the mammal, antibody molecules having immunospecificity for the immunizing peptide or peptides. 25 The peptide(s) or related molecule(s) may be monomeric, polymeric, conjugated to a carrier, and/or administered in the presence of an adjuvant. The antibody molecules may then be collected from the mammal if they are to be used in immunoassays or for providing passive immunity.

30 The antibody molecules of the present invention may be polyclonal or monoclonal. Monoclonal antibodies may be produced by methods known in the art. Portions of immunoglobulin molecules may also be produced by methods known in the art.

35

-21-

0 The antibody of the present invention may be
contained in various carriers or media, including blood,
plasma, serum (e.g., fractionated or unfractionated
serum), hybridoma supernatants and the like.
Alternatively, the antibody of the present invention is
isolated to the extent desired by well known techniques
5 such as, for example, by using DEAE Sephadex, or affinity
chromatography. The antibodies may be purified so as to
obtain specific classes or subclasses of antibody such as
IgM, IgG, IgA, IgG₁, IgG₂, IgG₃, IgG₄, and the like.
Antibody of the IgG class are preferred for purposes of
10 passive protection.

The presence of the antibodies of the present
invention, either polyclonal or monoclonal, can be
determined by various assays. Assay techniques include,
but are not limited to, immunobinding, immunofluorescence
15 (IF), indirect immunofluorescence, immunoprecipitation,
ELISA, agglutination and Western blot techniques.

The antibodies of the present invention have a number
of diagnostic and therapeutic uses. The antibodies can be
used as an in vitro diagnostic agent to test for the
20 presence of various staphylococcal and streptococcal
pyrogenic exotoxins in biological samples in standard
immunoassay protocols and to aid in the diagnosis of
various diseases related to the presence of bacterial
pyrogenic exotoxins. Preferably, the assays which use the
25 antibodies to detect the presence of bacterial pyrogenic
exotoxins in a sample involve contacting the sample with
at least one of the antibodies under conditions which will
allow the formation of an immunological complex between
the antibody and the toxin that may be present in the
30 sample. The formation of an immunological complex if any,
indicating the presence of the toxin in the sample, is
then detected and measured by suitable means. Such assays
include, but are not limited to, radioimmunoassays, (RIA),
ELISA, indirect immunofluorescence assay, Western blot and
35

-22-

the like. The antibodies may be labeled or unlabeled depending on the type of assay used. Labels which may be coupled to the antibodies include those known in the art and include, but are not limited to, enzymes, radionucleotides, fluorogenic and chromogenic substrates, cofactors, biotin/avidin, colloidal gold and magnetic particles. Modification of the antibodies allows for coupling by any known means to carrier proteins or peptides or to known supports, for example, polystyrene or polyvinyl microliter plates, glass tubes or glass beads and chromatographic supports, such as paper, cellulose and cellulose derivatives, and silica.

Such assays may be, for example, of direct format (where the labelled first antibody reacts with the antigen), an indirect format (where a labelled second antibody reacts with the first antibody), a competitive format (such as the addition of a labelled antigen), or a sandwich format (where both labelled and unlabelled antibody are utilized), as well as other formats described in the art. In one such assay, the biological sample is contacted to antibodies of the present invention and a labelled second antibody is used to detect the presence of staphylococcal and streptococcal pyrogenic exotoxins, to which the antibodies are bound.

The antibodies of the present invention are also useful as therapeutic agents in the prevention and treatment of diseases caused by the deleterious effects of staphylococcal and streptococcal pyrogenic exotoxins.

The antibodies are generally administered with a physiologically acceptable carrier or vehicle therefor. A physiologically acceptable carrier is one that does not cause an adverse physical reaction upon administration and one in which the antibodies are sufficiently soluble and retain their activity to deliver a therapeutically effective amount of the compound. The therapeutically effective amount and method of administration of the

-23-

antibodies may vary based on the individual patient, the indication being treated and other criteria evident to one of ordinary skill in the art. A therapeutically effective amount of the antibodies is one sufficient to attenuate the dysfunction without causing significant side effects such as non-specific T cell lysis or organ damage. The route(s) of administration useful in a particular application are apparent to one of ordinary skill in the art.

Routes of administration of the antibodies include, but are not limited to, parenteral, and direct injection into an affected site. Parenteral routes of administration include but are not limited to intravenous, intramuscular, intraperitoneal and subcutaneous.

The present invention includes compositions of the antibodies described above, suitable for parenteral administration including, but not limited to, pharmaceutically acceptable sterile isotonic solutions. Such solutions include, but are not limited to, saline and phosphate buffered saline for intravenous, intramuscular, intraperitoneal, subcutaneous or direct injection into a joint or other area.

Antibodies for use to elicit passive immunity in humans are preferably obtained from other humans previously inoculated with compositions comprising one or more of the consensus amino acid sequences of the invention. Alternatively, antibodies derived from other species may also be used. Such antibodies used in therapeutics suffer from several drawbacks such as a limited half-life and propensity to elicit an immune response. Several methods have been proposed to overcome these drawbacks. Antibodies made by these methods are encompassed by the present invention and are included herein. One such method is the "humanizing" of non-human antibodies by cloning the gene segment encoding the antigen binding region of the antibody to the human gene

-24-

segments encoding the remainder of the antibody. Only the binding region of the antibody is thus recognized as foreign and is much less likely to cause an immune response. An article describing such antibodies is Reichmann et al., "Reshaping Human Antibodies for Therapy", Nature 332:323-327 (1988), which is incorporated herein by reference.

In providing the antibodies of the present invention to a recipient mammal, preferably a human, the dosage of administered antibodies will vary depending upon such factors as the mammal's age, weight, height, sex, general medical condition, previous medical history and the like.

In general, it is desirable to provide the recipient with a dosage of antibodies which is in the range of from about 5 mg/kg to about 20 mg/kg body weight of the mammal, although a lower or higher dose may be administered. In general, the antibodies will be administered intravenously (IV) or intramuscularly (IM).

The antibodies of the present invention are intended to be provided to the recipient subject in an amount sufficient to prevent, or attenuate the severity, extent or duration of the deleterious effects of staphylococcal and streptococcal pyrogenic exotoxins.

The administration of the agents including peptide and antibody compositions of the invention may be for either "prophylactic" or "therapeutic" purpose. When provided prophylactically, the agents are provided in advance of any symptom. The prophylactic administration of the agent serves to prevent or ameliorate any subsequent deleterious effects of staphylococcal and streptococcal pyrogenic exotoxins. When provided therapeutically, the agent is provided at (or shortly after) the onset of a symptom of infection with bacteria expressing staphylococcal or streptococcal pyrogenic exotoxins. The agent of the present invention may, thus, be provided either prior to the anticipated exposure to

35

-25-

- o bacteria expressing staphylococcal or streptococcal pyrogenic exotoxin (so as to attenuate the anticipated severity, duration or extent of disease symptoms) or after the initiation of the infection.

Also envisioned are therapies based upon vectors, such as viral vectors containing nucleic acid sequences coding for the peptides described herein. These molecules, developed so that they do not provoke a pathological effect, will stimulate the immune system to respond to the peptides.

For all therapeutic, prophylactic and diagnostic uses, the peptide of the invention, alone or linked to a carrier, as well as antibodies and other necessary reagents and appropriate devices and accessories may be provided in kit form so as to be readily available and easily used.

Where immunoassays are involved, such kits may contain a solid support, such as a membrane (e.g., nitrocellulose), a bead, sphere, test tube, rod, and so forth, to which a receptor such as an antibody specific for the target molecule will bind. Such kits can also include a second receptor, such as a labelled antibody. Such kits can be used for sandwich assays to detect toxins. Kits for competitive assays are also envisioned.

The following examples illustrate certain embodiments of the present invention, but should not be construed as limiting its scope in any way. Certain modifications and variations will be apparent to those skilled in the art from the teachings of the foregoing disclosure and the following examples, and these are intended to be encompassed by the spirit and scope of the invention.

EXAMPLE 1

Peptides whose sequences are based on the two highly conserved regions of the staphylococcal and streptococcal pyrogenic exotoxins described herein were constructed. The sequences were based on alignments of the

-26-

streptococcal pyrogenic exotoxins with the staphylococcal enterotoxins, and the amino acids used in positions with possible degeneracy were the amino acids most frequently found in these positions. Three of the peptides were then catenated and polymerized to produce peptides of greater than 8000 daltons (i.e., peptides 6343, 6345 and 6348, described below). As described further below, peptide 6348 was used to immunize rabbits, which produced high titer antibodies to this peptide. These antibodies were tested for the ability to recognize the streptococcal and staphylococcal pyrogenic exotoxins. Immunological assays (immunoblots) revealed that these antibodies recognized regions common to all the pyrogenic exotoxins. These antibodies were also tested for the ability to neutralize in vitro and in vivo biological activity of the pyrogenic exotoxins. These antibodies protected against the biological T-cell proliferation of these toxins in an in vitro blastogenesis assay using human mononuclear cell populations. The lethal effects of staphylococcal toxin SEB and streptococcal pyrogenic toxin SPEA in vivo were also completely blocked by mixing the antibodies with the toxin prior to injection.

Materials and Methods

Construction of Synthetic Peptides:

Peptides were constructed by solid phase synthesis (20) using the modifications described by Houghton (10).

- | | | |
|----|------------------|--|
| 1. | GCG Consensus #1 | YGGX ₁ TX ₂ X ₃ X ₄ X ₅ N (SEQ ID NO:1) |
| | peptide #1 | CMYGGVTEHEGN (SEQ ID NO:3) |
| 2. | GCG Consensus #2 | KX ₆ X ₇ X ₈ X ₉ X ₁₀ X ₁₁ X ₁₂ X ₁₃ DX ₁₄ X ₁₅ X ₁₆ RX ₁₇ X ₁₈ L |
| | | X ₁₉ X ₂₀ X ₂₁ X ₂₂ X ₂₃ X ₂₄ Y (SEQ ID NO:2) |
| | peptide #2 | KKNVTVQELDYKIRKYLVDNKKLY (SEQ ID NO:4) |

As is evident above, synthetic peptides #1 and

-27-

5 #2 are not native peptides, i.e., their sequences differ from those found in native toxins. Variations of these peptides have also been constructed in order to generate concatenated polymers of the peptides. These polymers were constructed by the addition of glycine and of additional cysteine residues to the amino- and/or carboxyl- termini of the initial 2 peptides, thus facilitating concatenation via disulfide bond formation (37, 38, 39). The polymerized molecules were then dialyzed to remove molecules with molecular weights less than 6000-8000 daltons. One polymeric construct is composed of the monomer: CMYGGVTEHEGNGC (SEQ ID NO:5). An additional polymer is composed of the peptide: CGKKNVTVQELDYKIRKYLVDNKKLYGC (SEQ ID NO:6).

15 In the native toxin molecules, consensus region #1 precedes consensus region #2 by 27 amino acid residues (e.g. [consensus region 1] x27 [consensus region 2]). We have constructed the peptide: CMYGGVTEHEGNKKNVTVQELDYKIRKYLVDNKKLY (SEQ ID NO:7). Like the native toxin molecule, this peptide is representative of the two consensus regions joined together in the proper order (region 1 in the N terminal half, and region 2 in the C-terminal half of the molecule), however they are not separated by an additional 27 residues as they are in the native toxins. We have also constructed concatenated polymers based on the monomer: CMYGGVTEHEGNKKNVTVQELDYKIRKYLVDNKKLYGC (SEQ ID NO:8).

ID#	Peptide
6343	CMYGGVTEHEGN (SEQ ID NO:3)
6344	CMYGGVTEHEGNGC* (SEQ ID NO:5)
6345	KKNVTVQELDYKIRKYLVDNKKLY (SEQ ID NO:4)
30 6346	CGKKNVTVQELDYKIRKYLVDNKKLYGC* (SEQ ID NO:6)
6347	CMYGGVTEHEGNKKNVTVQELDYKIRKYLVDNKKLY (SEQ ID NO:7)
6348	CMYGGVTEHEGNKKNVTVQELDYKIRKYLVDNKKLYGC* (SEQ ID NO:8)

Peptides with an (*) are cross-linked polymers composed of the described sequence. It is expected that monomers of

35

-28-

these peptides will also be useful in the present invention.

Generation of anti-peptide sera.

New Zealand White rabbits were immunized by subcutaneous injection with 500 μ g of peptide in complete Freund's adjuvant. Additional booster injections of 500 μ g in incomplete adjuvant was administered 4 weeks after the primary injections. Ten days after booster injections, the rabbits were bled, and the anti peptide titers were determined by ELISA.

Staphylococcal enterotoxins, TSST-1, and streptococcal pyrogenic exotoxins were purchased from Toxin Technology Inc. (Sarasota, FL).

Immunoblots

Each of the staphylococcal and streptococcal pyrogenic exotoxins were electrophoresed through 10% SDS PAGE gels (16) and transferred to nitrocellulose for western blots (33). The western blots were developed using the rabbit anti-peptide 6348 serum (anti-pep 6348 or AP6348) diluted 1:5000, followed by goat anti-rabbit (IgG) alkaline phosphate conjugate (Sigma).

Inhibition of blastogenesis

Human peripheral blood mononuclear cell (PBMC) preparations were stimulated by each of the staphylococcal enterotoxins and streptococcal pyrogenic exotoxins. 100 ng of toxin was used to stimulate PBMC preparations at cell concentrations of 10^5 cells per well in 96 well microliter plates. Phytohemagglutinin (PHA) was used in place of the toxins as a positive mitogenic control. Cell culture medium was supplemented with either 10% normal rabbit serum (NRS) or AP6348 serum. Blastogenesis was assayed by incorporation of tritiated thymidine after 5 days of culture (22). All experiments were performed in triplicate.

Passive protection of rabbits

Female New Zealand White rabbits >1yr old were

-29-

obtained from Hazelton Dutchland Labs, Inc. (Denver, PA). Rabbits were challenged with staphylococcal or streptococcal toxins at doses ranging from 50 to 100 $\mu\text{g/kg}$, as previously described (24). Briefly, pyrogenic toxins were incubated with normal rabbit serum or anti-pep #6348 serum for one hour prior to challenge. Toxin-serum mixtures were administered intravenously through the marginal ear veins. Normal control rabbits were treated in an identical manner, with isotonic saline substituted for the pyrogenic toxin. Four hours later, rabbits were given a sub-lethal dose (5 $\mu\text{g/kg}$) of endotoxin (E. coli LPS, List Biological Laboratories, Inc., Campbell, CA). Rabbits were monitored 72 h for clinical signs of toxic shock. These included elevated temperature, diarrhea, cardiopulmonary distress, and conjunctival injection. Rabbits with severe toxic shock exhibiting cyanosis and temperatures less than 97°F were declared moribund. Moribund rabbits were euthanized by administration of 5 ml pentobarbital sodium. All animal protocols were reviewed by the Laboratory Animal Research Center at the Rockefeller University.

Results

ELISA assays

As seen in Figure 2, rabbits raised significant antibody titers to peptide 6348. Similarly, rabbits receiving immunizations with peptides 6344 and 6346 also developed high titers.

Recognition of staphylococcal and streptococcal toxins by anti-pep 6348 serum

Western blots of the staphylococcal and streptococcal toxins were developed with anti-peptide 6348 serum followed by an anti-rabbit IgG alkaline phosphatase conjugate (Sigma). The results indicate the anti-peptide 6348 serum recognizes the conserved regions of the bacterial toxin molecules; SEA, SEB, SEC, SEE, SPEA, SPEC, and TSST-1 (Figure 3). SED did not show a significant

-30-

reaction with anti-peptide 6348.

Blastogenesis inhibition

The percentage of inhibition, of toxin mediated blastogenesis, by AP6348 was assayed. Tritiated thymidine incorporation by human PBMC stimulated with staphylococcal and streptococcal pyrogenic toxins was significantly inhibited by the addition of AP6348 compared to normal rabbit serum (NRS) (Figure 4). This suggests blastogenesis of PBMC in response to the toxins was inhibited by AP6348. The AP6348 serum did not affect the blastogenesis of human PBMC in response to PHA, suggesting a specific inhibition of toxin biologic activity.

In vivo protection of rabbits

We tested the ability of AP6348 serum to prevent severe toxic shock in rabbits challenged with SEB and NRS. Rabbits challenged intravenously with a mixture of SEB and NRS developed symptoms of severe toxic shock (Table 2). One rabbit receiving 50 $\mu\text{g/kg}$ SEB with NRS, and two receiving 100 $\mu\text{g/kg}$ of SEB with NRS, developed severe toxic shock and were declared moribund within 30 hrs. In contrast, two rabbits challenged with 50 $\mu\text{g/kg}$ and 100 $\mu\text{g/kg}$ SEB with AP6348 developed fever, but this returned to normal by 32 hours. No diarrhea or cardiopulmonary depression was observed. Rabbits were followed for a total of 5 days (data not shown) and appeared fully recovered.

-31-

Table 2Passive Protection of Rabbits Challenged with SEB, SPEA and LPS

	<u>Toxin</u>	<u>LPS</u>	<u>Diarrhea</u>	<u>Temperature °F</u>				
				0 hr	4 hr	24 hr	32 hr	48 hr
5	<u>SEB</u>							
	ns\NRS	5	-	100.4	102	101.4	101.2	NT
	50\NRS	5	+	101	104.4	102.8	96◇	
	100\NRS	5	+	102	104.6	103	97◇	
10	100\NRS	5	+	101	104.5	102.6	97◇	
	50\APS	5	-	101.4	103.8	103	102	101
	100\APS	5	-	100.4	104.4	103	102	101
15	<u>SPEA</u>							
	50\NRS	5	+	101	104.2	NT◇		
	100\NRS	5	+	102	104.8	NT◇		
	50\APS	5	-	102	104	103	102	102
20	100\APS	5	+	101.6	104.4	104	100	97◇

ns^f=control rabbit given isotonic saline in place of SEB or SPEA

NRS=Normal rabbit serum

APS=Anti-peptide 6348 serum

◇=animals were declared moribund

NT=not taken

Discussion

Our results demonstrate that antibodies rabbit antiserum generated to peptides representative of two regions with highly conserved amino acid sequences (AP6348) are capable of recognizing most of the staphylococcal enterotoxins and streptococcal pyrogenic exotoxins (e.g. SEA, SEB, SEC, SEE, SPEA, SPEC), as well

-32-

as TSST-1, using Western blots. We expect that other, more sensitive assays, will result in the demonstration of binding of these antibodies to additional members, probably all members, of the staphylococcal and streptococcal pyrogenic toxin family.

Since recognition of the toxins by AP6348 was successful, we tested this serum for the ability to inhibit the biological effects of these pyrogenic toxins. AP6348 was capable of inhibiting in vitro blastogenesis of human PBMCs by many of the pyrogenic toxins (e.g., SEA, SEB, SEC, SEE, SPEA, and SPEC).

AP6348 was also able to provide passive in vivo protection of animals challenged with lethal doses of SEB and SPEA. These animals developed fever, however the fever returned to normal within 30 hours and remained normal. Rabbits appeared to be fully recovered within days of challenge.

In contrast, rabbits receiving similar doses of SEB and SPEA pre-incubated with NRS developed severe toxic shock, as evidenced by high fevers, diarrhea, and cardiopulmonary distress. The illness progressed and these animals were declared moribund.

The therapeutic and biological implications of these observations are as follows: (i) antibodies prepared against this peptide may be administered during the early stages of toxic shock irrespective of the toxin causing the symptoms and (ii) the peptide may be used as an immunogen to block the toxic effects of this family of superantigens.

-33-

°
References

1. Bergdoll, M. S. 1985. The staphylococcal enterotoxins-an update., 247-254. In J. Jeljaszewicz (ed.). The staphylococci. Gustav Fischer Verlag, New York, NY.
- 5 2. Blomster-Hautamaa, D. A., B. N. Kreiswirth, J. S. Kornblum, R. P. Novick and P. M. Schlievert. 1986. The nucleotide and partial amino acid sequence of toxic shock syndrome toxin-1. Journal of Biological Chemistry. 261:15783-15786.
- 10 3. Choi, Y., B. Kotzin, L. Herron, J. Callahan, P. Marrack and J. Kappler. 1989. Interaction of staphylococcus aureus toxin superantigens with human T cells. Proc. Natl. Acad. Sci. USA. 86:8941.
- 15 4. Fleischer, B. and H. Schrezenmeier. 1988. T cell stimulation by staphylococcal enterotoxins. Clonally variable response and requirement for major histocompatibility complex class II molecules on accessory or target cells. Journal of Experimental Medicine. 167:1697.
- 20 5. Goshorn, S. C. and P. M. Schlievert. 1988. Nucleotide sequence of streptococcal exotoxin type C. Infect. Immun. 56:2518-2520.
- 25 6. Grossman, D., M. Van, J. A. Mollick, S. K. Highlander and R. R. Rich. 1991. Mutation of the disulfide loop in staphylococcal enterotoxin A. Consequences for T cell recognition. J. Immunol. 147:3274-3281.
- 30 7. Hartwig, U. F. and B. Fleisher. 1993. Mutations affecting MHC class II binding of the superantigen streptococcal erythrogenic toxin A. International Immunology. 5:869-875.
- 35 8. Hauser, A. R., D. L. Stevens, E. L. Kaplan and P. M. Schlievert. 1991. Molecular analysis of pyrogenic exotoxins from streptococcus pyogenes isolates

-34-

o associated with toxic shock-like syndrome. Journal of Clinical Microbiology. 29:1562-1567.

9. Hensler, T., M. Köller, C. Geoffroy, J. E. Alouf and W. König. 1993. Staphylococcus aureus toxic shock syndrome toxin 1 and streptococcus pyogenes erythrogenic toxin A modulate inflammatory mediator
5 release from human neutrophils. Infect. Immun. 61:1055-1061.

10. Houghten, R.A. 1985. General method for the rapid solid phase synthesis of large numbers of peptides: specificity of antigen-antibody interactions at the level
10 of individual amino acids. Proc. Natl. Acad. Sci. USA. 82:5131-5135.

11. Hynes, W. L., C. R. Weeks, J. J. Iandolo and J. J. Ferretti. 1987. Immunologic cross-reactivity of type A streptococcal exotoxin (erythrogenic toxin) and
15 staphylococcal enterotoxins B and C1. Infect. Immun. 55:837-840.

12. Janeway, C. A. J., J. Yagi, P. J. Conrad, M. E. Katz, B. Jones, S. Vroegop and S. Buxser. 1989. T-cell responses to Mls and to bacterial proteins that
20 mimic its behavior. Immunology Reviews. 107:61.

13. Johnson, L. P., J. J. L'Italien and P. M. Schlievert. 1986. Streptococcal pyrogenic exotoxin type A (scarlet fever toxin) is related to staphylococcus aureus enterotoxin B. Molecular and General Genetics.
25 203:354-356.

14. Kappler, J., B. L. Kotzin, L. Herron, E. W. Gelfand, R. D. Bigler, A. Boylston, S. Carrell, D. N. Posnett, Y. Choi and P. Marrack. 1989. V β -specific stimulation of human T-cells by staphylococcal toxins.
30 Science. 248:705.

15. Kappler, J. W., A. Herman, J. Clements and P. Marrack. 1992. Mutations defining functional regions of the superantigen staphylococcal enterotoxin B. Journal of Experimental Medicine. 175:387-396.

35

-35-

16. Laemmli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*. 227:680.

17. Leonard, B. A. and P. M. Schlievert. 1992. Immune cell lethality induced by streptococcal pyrogenic exotoxin A and endotoxin. *Infect. Immun.* 60:3747-3755.

18. Marrack, P., M. Blackman, E. Kushnir and J. Kappler. 1990. The toxicity of staphylococcal enterotoxin B in mice is mediated by T cells. *Journal of Experimental Medicine*. 171:455.

19. Marrack, P. and J. Kappler. 1990. The staphylococcal enterotoxins and their relatives. *Science*. 248:705-711.

20. Merrifield, R.B. 1963. Solid-phase peptide synthesis I. The synthesis of a tetrapeptide. *J. Am. Chem. Soc.* 85:2149-2154.

21. Musser, J. M., A. R. Hauser, M. H. Kim, P. M. Schlievert, K. Nelson and R. K. Selander. 1991. *Streptococcus pyogenes* causing toxic-shock-like syndrome and other invasive diseases: clonal diversity and pyrogenic exotoxin expression. *Proceedings of the National Academy of Sciences of the United States of America*. 88:2668-2672.

22. Read, S. E., H. F. M. Reid, V. A. Fischetti, T. Poon-King, R. Ramkisson, M. McDowell and J. B. Zabriskie. 1986. Serial studies on the cellular immune response to streptococcal antigens in acute and convalescent rheumatic fever patients in Trinidad. *Journal of Clinical Immunology*. 6:433-441.

23. Reda, K. B., V. Kapur, J. A. Mollick, J. G. Lamphear, J. M. Musser and R. R. Rich. 1994. Molecular characterization and phylogenetic distribution of the streptococcal superantigen gene (ssa) from *streptococcus pyogenes*. *Infect. Immun.* 62:1867-1874.

24. Ren, K., J. D. Bannan, V. Pancholi, A. L. Cheung, J. C. Robbins, V. A. Fischetti and J. B.

-36-

- 0 Zabriskie. 1994. Characterization and biological
properties of a new staphylococcal exotoxin. *Journal of
Experimental Medicine*. 180:1675-1683.
- 5 25. Smith, R. J., P. M. Schlievert, I. M.
Himelright and L. M. Baddour. 1994. Dual infections with
staphylococcus aureus and streptococcus pyogenes causing
toxic shock syndrome. Possible synergistic effects of
toxic shock syndrome toxin 1 and streptococcal pyrogenic
exotoxin C. *Diagnostic Microbiology & Infectious Disease*.
19:245-247.
- 10 26. Spero, L., B. Morlock and J. Metzger.
1978. On the cross-reactivity of staphylococcal
enterotoxins A, B, and C. *J. Immunol.* 120:86-89.
- 15 27. Spero, L. and B. A. Morlock. 1978.
Biological activities of the peptides of staphylococcal
enterotoxin C formed by limited tryptic hydrolysis.
Journal of Biological Chemistry. 253:8787-8791.
- 20 28. Spero, L. and B. A. Morlock. 1979.
Cross-reactions between tryptic polypeptides of
staphylococcal enterotoxins B and C. *J. Immunol.*
122:1285-1289.
- 25 29. Stelma, G. N., Jr. and M. S. Bergdoll.
1982. Inactivation of staphylococcal enterotoxin A by
chemical modification. *Biochemical and Biophysical
Research Communications*. 105:121-126.
- 30 30. Stevens, D. L., M. H. Tanner, J. Winship,
R. Swarts, K. M. Ries, P. M. Schlievert and E. Kaplan.
1989. Severe group A streptococcal infections associated
with a toxic shock-like syndrome and scarlet fever toxin
A. *The New England Journal of Medicine*. 321:1-7.
- 35 31. Sugiyama, H., E. M. J. McKissic, M. S.
Bergdoll and B. Heller. 1964. Enhancement of bacterial
endotoxin lethality by staphylococcal enterotoxin. *J.
Infect. Dis.* 114:111-118.
32. Swaminathan, S., W. Furey, J. Pletcher and
M. Sax. 1992. Crystal structure of staphylococcal

-37-

- enterotoxin B, a superantigen. *Nature*. 359:801-806.
33. Towbin, H., T. Staehlin and J. Gordon. 1979. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets. Procedure and some applications. *Proceedings of the National Academy of Sciences, USA*. 76:4350.
34. Van den Bussche, R. A., J. D. Lyon and G. A. Bohac. 1993. Molecular evolution of the staphylococcal and streptococcal pyrogenic toxin gene family. *Molecular Phylogenetics and Evolution*. 2:281-292.
35. Weeks, C. R. and J. J. Ferretti. 1986. Nucleotide sequence of the type A streptococcal exotoxin (erythrogenic toxin) gene from streptococcus pyogenes bacteriophage T12. *Infect. Immun.* 52:144-150.
36. White, J., A. Herman, A. M. Pullen, R. Kubo, J. W. Kappler and P. Marrack. 1989. The V beta-specific superantigen staphylococcal enterotoxin B: stimulation of mature T cells and clonal deletion in neonatal mice. *Cell*. 56:27-35.
37. Patarroyo, M.E., R. Amador, P. Clavijo, A. Moreno, F. Guzman, P. Romero, R. Tascon, A. Franco, L.A. Murillo, G. Ponton and G. Trujillo. 1988. A synthetic vaccine protects humans against challenge with *Plasmodium falciparum* malaria. *Nature*. 332:158-161.
38. Lopez, M.C., Y. Silva, M.C. Thomas, A. Garcia, M.J. Faus, P. Alonso, F. Martinez, G. Del Real and C. Alonso. 1994. Characterization of SPf(66)n: a chimeric molecule used as a malaria vaccine. *Vaccine*. 12:585-591.
39. Rodriguez, R., A. Moreno, F. Guzman, M. Calvo and M.E. Patarroyo. 1990. Studies in owl monkeys leading to the development of a synthetic vaccine against the asexual blood stages of *Plasmodium falciparum*. *Am. J. Trop. Med. Hyg.* 43:339-354.
40. Hoffman, M.L., L.M. Jablonski, K.K. Crum, S.P. Hackett, Y.-I. Chi, C.V. Stauffacher, D.L. Stevens

-38-

- o and G.A. Bohach. 1994. Predictions of T-cell receptor and Major Histocompatibility Complex-binding sites on staphylococcal enterotoxin C1. Infection and Immunity. 62:3396-3407.

5

10

15

20

25

30

35

-39-

Modifications of the above described modes for carrying out the invention that are obvious to those of skill in the fields of immunology, protein chemistry, microbiology, medicine, and related fields are intended to be within the scope of the following claims.

Every reference cited hereinbefore is hereby incorporated by reference in its entirety.

The following table shows the correspondence between peptides in Figure 1 and their sequence identification numbers:

Table 3

Correspondence between Sequence
Identification Numbers and Peptides in Figure 1

<u>Figure 1</u>				<u>Sequence ID Nos.</u>	
<u>Region 1</u>					
15	PEP		CMYGGVTEHEGN	SEQ ID NO:3	
	SEA	130	CMYGGVTLHDNN 141	SEQ ID NO:9	
	SEB	140	CMYGGVTEHNGN 151	SEQ ID NO:10	
	SEC	137	CMYGGITKHEGN 148	SEQ ID NO:11	
	SED	131	CTYGGVTPHEGN 142	SEQ ID NO:12	
	SEE	130	CMYGGVTLHDNN 141	SEQ ID NO:13	
	SEH	116	CLYGGITL.NSE 126	SEQ ID NO:14	
20	SPEA	128	CIYGGVTNHEGN 139	SEQ ID NO:15	
	SPEC	112	YIYGGITPAQNN 123	SEQ ID NO:16	
	SSA	134	CMYGGVTEHHRN 145	SEQ ID NO:17	
<u>Region 2</u>					
	PEP		KKNVTVQELDYKIRKYLVDNKKLY	SEQ ID NO:4	
	SEA	171	KKNVTVQELDLQARRYLQEKYNLY 194	SEQ ID NO:18	
25	SEB	179	KKKVTAQELDYLTRHYLVKNKKLY 202	SEQ ID NO:19	
	SEC	178	KKSMTAQELDIKARNFLINKKNLY 201	SEQ ID NO:20	
	SED	172	KKNVTVQELDAQARRYLQKDLKLY 195	SEQ ID NO:21	
	SEE	171	KKEVTVQELDLQARHYLHGKFGLY 194	SEQ ID NO:22	
	SEH	151	KKNVTLQELDIKIRKILSDKYKIY 174	SEQ ID NO:23	
	SPEA	167	KKMVTAQELDYKVRKYLTDNKQLY 190	SEQ ID NO:24	
	SPEC	151	KDIVTFQEIDFKIRKLYMDNYKIY 174	SEQ ID NO:25	
	SSA	174	KKQVTVQELDCKTRKILVSRKNLY 197	SEQ ID NO:26	
30	TSST1	161	KKQLAISTLDFEIRHQLTQIHGLY 184	SEQ ID NO:27	

-40-

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

(A) NAME: THE ROCKEFELLER UNIVERSITY
(B) STREET: 1230 YORK AVENUE
(C) CITY: NEW YORK
(D) STATE OR PROVINCE: NEW YORK
(E) COUNTRY: UNITED STATES OF AMERICA
(F) POSTAL CODE: 10021-6399

(ii) TITLE OF INVENTION: PEPTIDES USEFUL FOR
REDUCING SYMPTOMS OF TOXIC SHOCK SYNDROME

(iii) NUMBER OF SEQUENCES: 31

(iv)^a CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: MORGAN & FINNEGAN
(B) STREET: 345 PARK AVENUE
(C) CITY: NEW YORK
(D) STATE: NEW YORK
(E) COUNTRY: USA
(F) ZIP: 10154

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: FLOPPY DISK
(B) COMPUTER: IBM PC COMPATIBLE
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: WORDPERFECT 5.1

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: TO BE ASSIGNED
(B) FILING DATE: 01 APRIL 1998
(C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: 08/838,413
(B) FILING DATE: 07 APRIL 1997
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: MORRY, MARY J.
(B) REGISTRATION NUMBER: 34,398
(C) REFERENCE/DOCKET NUMBER: 2016-4010PC

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (212)758-4800
(B) TELEFAX: (212)751-6849

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

-41-

- (A) LENGTH: 10
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN

(ii) MOLECULE TYPE: PEPTIDE

(xi) SEQUENCE DESCRIPTIONS:SEQ ID NO: 1:

5 Tyr Gly Gly Xaa Thr Xaa Xaa Xaa Xaa Asn
5 10

(3) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 24
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN

(ii) MOLECULE TYPE: PEPTIDE

(xi) SEQUENCE DESCRIPTIONS:SEQ ID NO: 2:

15 Lys Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Asp Xaa Xaa
5 10

Xaa Arg Xaa Xaa Leu Xaa Xaa Xaa Xaa Xaa Xaa Tyr
15 20

20

(4) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 12
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN

(ii) MOLECULE TYPE: PEPTIDE

(xi) SEQUENCE DESCRIPTIONS:SEQ ID NO: 3:

30 Cys Met Tyr Gly Gly Val Thr Glu His Glu Gly Asn
5 10

35

(5) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- 35 (A) LENGTH: 24
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN

-42-

(D) TOPOLOGY: UNKNOWN

(ii) MOLECULE TYPE: PEPTIDE

(xi) SEQUENCE DESCRIPTIONS:SEQ ID NO: 4:

Lys Lys Asn Val Thr Val Gln Glu Leu Asp Tyr Lys
5 10

5 Ile Arg Lys Tyr Leu Val Asp Asn Lys Lys Leu Tyr
15 20

(6) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

10 (A) LENGTH: 14
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN

(ii) MOLECULE TYPE: PEPTIDE

(xi) SEQUENCE DESCRIPTIONS:SEQ ID NO: 5:

15 Cys Met Tyr Gly Gly Val Thr Glu His Glu Gly Asn
5 10

Gly Cys

20 (7) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN

25 (ii) MOLECULE TYPE: PEPTIDE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

Cys Gly Lys Lys Asn Val Thr Val Gln Glu Leu Asp
5 10

30 Tyr Lys Ile Arg Lys Tyr Leu Val Asp Asn Lys Lys
15 20

Leu Tyr Gly Cys
25

35 (8) INFORMATION FOR SEQ ID NO: 7:

-43-

- 0 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 36
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN
- (ii) MOLECULE TYPE: PEPTIDE
- 5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:
Cys Met Tyr Gly Gly Val Thr Glu His Glu Gly Asn
5 10
Lys Lys Asn Val Thr Val Gln Glu Leu Asp Tyr Lys
15 20
10 Ile Arg Lys Tyr Leu Val Asp Asn Lys Lys Leu Tyr
25 30 35
- (9) INFORMATION FOR SEQ ID NO: 8:
- 15 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 38
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN
- (ii) MOLECULE TYPE: PEPTIDE
- 20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:
Cys Met Tyr Gly Gly Val Thr Glu His Glu Gly Asn
5 10
Lys Lys Asn Val Thr Val Gln Glu Leu Asp Tyr Lys
15 20
25 Ile Arg Lys Tyr Leu Val Asp Asn Lys Lys Leu Tyr
25 30 35
Gly Cys
- (10) INFORMATION FOR SEQ ID NO: 9:
- 30 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN
- (ii) MOLECULE TYPE: PEPTIDE
- 35

-44-

o (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:
Cys Met Tyr Gly Gly Val Thr Leu His Asp Asn Asn
5 10

(11) INFORMATION FOR SEQ ID NO: 10:

5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN

(ii) MOLECULE TYPE: PEPTIDE

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:
Cys Met Tyr Gly Gly Val Thr Glu His Asn Gly Asn
5 10

(12) INFORMATION FOR SEQ ID NO: 11:

15 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN

(ii) MOLECULE TYPE: PEPTIDE

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:
Cys Met Tyr Gly Gly Ile Thr Lys His Glu Gly Asn
5 10

25 (13) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN

30 (ii) MOLECULE TYPE: PEPTIDE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:
Cys Thr Tyr Gly Gly Val Thr Pro His Glu Gly Asn
5 10

35

-45-

0 (14) INFORMATION FOR SEQ ID NO: 13:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN

5 (ii) MOLECULE TYPE: PEPTIDE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

Cys Met Tyr Gly Gly Val Thr Leu His Asp Asn Asn
5 10

10 (15) INFORMATION FOR SEQ ID NO: 14:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 11
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN

15 (ii) MOLECULE TYPE: PEPTIDE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

Cys Leu Tyr Gly Gly Ile Thr Leu Asn Ser Glu
5 10

20 (16) INFORMATION FOR SEQ ID NO: 15:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN

25 (ii) MOLECULE TYPE: PEPTIDE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

Cys Ile Tyr Gly Gly Val Thr Asn His Glu Gly Asn
5 10

30

(17) INFORMATION FOR SEQ ID NO: 16:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN

35

-46-

(D) TOPOLOGY: UNKNOWN

(ii) MOLECULE TYPE: PEPTIDE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

Tyr Ile Tyr Gly Gly Ile Thr Pro Ala Gln Asn Asn
5 10

(18) INFORMATION FOR SEQ ID NO: 17:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 12
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN

(ii) MOLECULE TYPE: PEPTIDE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

Cys Met Tyr Gly Gly Val Thr Glu His His Arg Asn
5 10

(19) INFORMATION FOR SEQ ID NO: 18:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN

(ii) MOLECULE TYPE: PEPTIDE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

Lys Lys Asn Val Thr Val Gln Glu Leu Asp Leu Gln
5 10

Ala Arg Arg Tyr Leu Gln Glu Lys Tyr Asn Leu Tyr
15 20

(20) INFORMATION FOR SEQ ID NO: 19:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN

(ii) MOLECULE TYPE: PEPTIDE

-47-

o (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

Lys Lys Lys Val Thr Ala Gln Glu Leu Asp Tyr Leu
5 10

Thr Arg His Tyr Leu Val Lys Asn Lys Lys Leu Tyr
15 20

5 (21) INFORMATION FOR SEQ ID NO: 20:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN

10

(ii) MOLECULE TYPE: PEPTIDE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

Lys Lys Ser Val Thr Ala Gln Glu Leu Asp Ile Lys
5 10

15

Ala Arg Asn Phe Leu Ile Asn Lys Lys Asn Leu Tyr
15 20

(22) INFORMATION FOR SEQ ID NO: 21:

20 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN

(ii) MOLECULE TYPE: PEPTIDE

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:

Lys Lys Asn Val Thr Val Gln Glu Leu Asp Ala Gln
5 10

Ala Arg Arg Tyr Leu Gln Lys Asp Leu Lys Leu Tyr
15 20

30

(23) INFORMATION FOR SEQ ID NO: 22:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH:
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN

35

-48-

o (D) TOPOLOGY: UNKNOWN

(ii) MOLECULE TYPE: PEPTIDE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:

Lys Lys Glu Val Thr Val Gln Glu Leu Asp Leu Gln
5 10

5 Ala Arg His Tyr Leu His Gly Lys Phe Gly Leu Tyr
15 20

(24) INFORMATION FOR SEQ ID NO: 23:

10 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN

(ii) MOLECULE TYPE: PEPTIDE

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:

Lys Lys Asn Val Thr Leu Gln Glu Leu Asp Ile Lys
5 10

Ile Arg Lys Ile Leu Ser Asp Lys Tyr Lys Ile Tyr
15 20

20 (25) INFORMATION FOR SEQ ID NO: 24:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN

25 (ii) MOLECULE TYPE: PEPTIDE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24:

Lys Lys Met Val Thr Ala Gln Glu Leu Asp Tyr Lys
5 10

30 Val Arg Lys Tyr Leu Thr Asp Asn Lys Gln Leu Tyr
15 20

(26) INFORMATION FOR SEQ ID NO: 25:

35 (i) SEQUENCE CHARACTERISTICS:

-49-

- o (A) LENGTH: 24
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN

(ii) MOLECULE TYPE: PEPTIDE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:

5 Lys Asp Ile Val Thr Phe Gln Glu Ile Asp Phe Lys
5 10
Ile Arg Lys Leu Tyr Met Asp Asn Tyr Lys Ile Tyr
15 20

10 (27) INFORMATION FOR SEQ ID NO: 26:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 24
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN

15 (ii) MOLECULE TYPE: PEPTIDE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:

Lys Lys Gln Val Thr Val Gln Glu Leu Asp Cys Lys
5 10
20 Thr Arg Lys Ile Leu Val Ser Arg Lys Asn Leu Tyr
15 20

(28) INFORMATION FOR SEQ ID NO: 27:

- 25 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 24
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN

(ii) MOLECULE TYPE: PEPTIDE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 27:

30 Lys Lys Gln Leu Ala Ile Ser Thr Leu Asp Phe Glu
5 10
Ile Arg His Gln Leu Thr Gln Ile His Gly Leu Tyr
15 20

35

-50-

0 (29) INFORMATION FOR SEQ ID NO: 28:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN

5 (ii) MOLECULE TYPE: PEPTIDE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 28:

Xaa Xaa Tyr Gly Gly Xaa Thr Xaa Xaa Xaa Xaa Asn
5 10

10 (30) INFORMATION FOR SEQ ID NO: 29:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 24
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN

15 (ii) MOLECULE TYPE: PEPTIDE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 29:

Lys Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Asp Xaa Xaa
5 10

20 Xaa Arg Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Tyr
15 20

(31) INFORMATION FOR SEQ ID NO: 30:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12
(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN

25 (ii) MOLECULE TYPE: PEPTIDE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 30:

30 Cys Met Tyr Gly Gly Xaa Thr Xaa His Xaa Gly Asn
5 10

(32) INFORMATION FOR SEQ ID NO: 31:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 24

35

-51-

(B) TYPE: AMINO ACID
(C) STRANDEDNESS: UNKNOWN
(D) TOPOLOGY: UNKNOWN

(ii) MOLECULE TYPE: PEPTIDE

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 31:

5 Lys Lys Xaa Val Thr Xaa Gln Glu Leu Asp Xaa Xaa
5 10

Xaa Arg Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Leu Tyr
15 20

10

15

20

25

30

35

-52-

We claim:

1. A peptide comprising a consensus amino acid sequence selected from the group consisting of $X_{25}X_{26}YGGX_1TX_2X_3X_4X_5N$ (SEQ ID NO:28) and $KX_6X_7X_8X_9X_{10}X_{11}X_{12}X_{13}DX_{14}X_{15}X_{16}RX_{17}X_{18}X_{27}X_{19}X_{20}X_{21}X_{22}X_{23}X_{24}Y$ (SEQ ID NO:29) wherein X_1 , X_8 , X_{13} and X_{24} are each independently selected from the group consisting of L, I and V; X_2 , X_4 , X_5 , X_6 , X_7 , X_9 , X_{10} , X_{11} , X_{12} , X_{14} , X_{15} , X_{16} , X_{17} , X_{18} , X_{19} , X_{20} , X_{21} , X_{22} , and X_{23} are each independently selected from the group consisting of any amino acid; X_3 , X_{25} and X_{26} are each independently selected from the group consisting of any amino acid and of no amino acid; and X_{27} is selected from the group consisting of L and Y.
2. A peptide of claim 1 wherein
 - X_2 is selected from the group consisting of L, E, K, P and N;
 - X_3 is selected from the group consisting of H and A or no amino acid;
 - X_4 is selected from the group consisting of D, N, E, Q, and H;
 - X_5 is selected from the group consisting of N, G, S, and R;
 - X_6 is selected from the group consisting of K and D;
 - X_7 is selected from the group consisting of N, K, S, E, M, I and Q;
 - X_9 is selected from the group consisting of T and A;
 - X_{10} is selected from the group consisting of V, A, L, F and I;
 - X_{11} is selected from the group consisting of Q and S;
 - X_{12} is selected from the group consisting of E and T;
 - X_{14} is selected from the group consisting of L, Y, I, A, F and C;

-53-

0 X₁₅ is selected from the group consisting of Q,
L, K and E;

 X₁₆ is selected from the group consisting of A,
T, I and V;

 X₁₇ is selected from the group consisting of R,
H, N and K;

5 X₁₈ is selected from the group consisting of Y,
F, I, L and Q;

 X₁₉ is selected from the group consisting of Q,
V, I, H, S, T and M;

10 X₂₀ is selected from the group consisting of E,
K, N, G, D, S and Q;

 X₂₁ is selected from the group consisting of K,
N, D, R and I;

 X₂₂ is selected from the group consisting of Y,
K, L, F and H;

15 X₂₃ is selected from the group consisting of N,
K, G and Q;

 X₂₅ is selected from the group consisting of C,
Y and no amino acid; and

20 X₂₆ is selected from the group consisting of M,
T, L, I and no amino acid.

3. A peptide of claim 1 wherein X₁ is
valine; X₂ is leucine; X₃ is histidine; X₄ is glutamic
acid; X₅ is glycine; X₆ is lysine; X₇ is asparagine; X₈ is
valine; X₉ is threonine; X₁₀ is valine; X₁₁ is glutamine;
25 X₁₂ is glutamic acid; X₁₃ is leucine; X₁₄ is leucine,
tyrosine, isoleucine or phenylalanine; X₁₅ is lysine; X₁₆
is alanine or isoleucine; X₁₇ is lysine; X₁₈ is tyrosine;
X₁₉ is glutamine, valine or threonine; X₂₀ is aspartic
acid; X₂₁ is lysine; X₂₂ is lysine; X₂₃ is lysine; X₂₄ is
30 leucine; X₂₅ is cysteine; X₂₆ is methionine and X₂₇ is
leucine.

4. A peptide comprising at least one amino
acid sequence selected from the group consisting of
CMYGGVTEHEGN (SEQ ID NO: 3), CMYGGVTEHEGNGC (SEQ ID NO:

35

-54-

5), KKNVTVQELDYKIRKYLVDNKKLY (SEQ ID NO: 4),
 CGKKNVTVQELDYKIRKYLVDNKKLYGC (SEQ ID NO: 6),
 CMYGGVTEHEGNKKNVTVQELDYKIRKYLVDNKKLY (SEQ ID NO: 7), and
 CMYGGVTEHEGNKKNVTVQELDYKIRKYLVDNKKLYGC (SEQ ID NO: 8).

5. A peptide of claim 4 wherein the peptide
 comprises the amino acid sequence

CMYGGVTEHEGNKKNVTVQELDYKIRKYLVDNKKLYGC (SEQ ID NO: 8).

6. A peptide of claim 4 wherein the amino acid
 sequence is a component of a larger molecule of at least
 6,000 to 8,000 daltons.

7. A pharmaceutical composition
 comprising a peptide comprising a consensus amino acid
 sequence selected from the group consisting of
 $X_{25}X_{26}YGGX_1TX_2X_3X_4X_5N$ (SEQ ID NO:28) and
 $KX_6X_7X_8X_9X_{10}X_{11}X_{12}X_{13}DX_{14}X_{15}X_{16}RX_{17}X_{18}X_{27}X_{19}X_{20}X_{21}X_{22}X_{23}X_{24}Y$ (SEQ ID
 NO:29) wherein X_1 , X_8 , X_{13} and X_{24} are each independently
 selected from the group consisting of L, I and V; X_2 , X_4 ,
 X_5 , X_6 , X_7 , X_9 , X_{10} , X_{11} , X_{12} , X_{14} , X_{15} , X_{16} , X_{17} , X_{18} , X_{19} , X_{20} ,
 X_{21} , X_{22} , and X_{23} are each independently selected from the
 group consisting of any amino acid; X_3 , X_{25} and X_{26} are each
 independently selected from the group consisting of any
 amino acid and of no amino acid; and X_{27} is selected from
 the group consisting of L and Y;
 in a physiologically acceptable carrier.

8. A pharmaceutical composition comprising a
 peptide of claim 4 in a physiologically acceptable
 carrier.

9. A method of inducing serum antibodies
 that bind at least one staphylococcal enterotoxin or
 streptococcal exotoxin, said method comprising
 administering to a mammal, in a physiologically acceptable
 carrier, an amount of a peptide comprising a consensus
 amino acid sequence selected from the group consisting of
 $X_{25}X_{26}YGGX_1TX_2X_3X_4X_5N$ (SEQ ID NO:28) and
 $KX_6X_7X_8X_9X_{10}X_{11}X_{12}X_{13}DX_{14}X_{15}X_{16}RX_{17}X_{18}X_{27}X_{19}X_{20}X_{21}X_{22}X_{23}X_{24}Y$ (SEQ ID NO:
 29) wherein X_1 , X_8 , X_{13} and X_{24} are each independently

-55-

selected from the group consisting of L, I and V; X₂, X₄, X₅, X₆, X₇, X₉, X₁₀, X₁₁, X₁₂, X₁₄, X₁₅, X₁₆, X₁₇, X₁₈, X₁₉, X₂₀, X₂₁, X₂₂, and X₂₃ are each independently selected from the group consisting of any amino acid; X₃, X₂₅ and X₂₆ are each independently selected from the group consisting of any amino acid and of no amino acid; and X₂₇ is selected from the group consisting of L and Y; sufficient to elicit production of said antibodies.

10. The method of claim 9 wherein said peptide is administered at a dose of about 5 micrograms to about 100 micrograms.

11. A method of inducing serum antibodies that bind at least one staphylococcal enterotoxin or streptococcal exotoxin, said method comprising administering to a mammal, in a physiologically acceptable carrier, an amount of a peptide of claim 4 sufficient to elicit production of said antibodies.

12. The method of claim 11 wherein the peptide comprises the amino acid sequence
CMYGGVTEHEGNKKNVTVQELDYKIRKYLVDNKKLYGC (SEQ ID NO: 8).

13. The method of claim 11 wherein said peptide is administered at a dose of about 5 micrograms to about 100 micrograms.

14. A method of inducing serum antibodies which detect the toxins SPEA, SEA, SEB, and SED, comprising administering to a mammal, in a physiologically acceptable carrier, an immunologically sufficient amount of a peptide comprising a consensus amino acid sequence selected from the group consisting of X₂₅X₂₆YGGX₁TX₂X₃X₄X₅N (SEQ ID NO:28) and KX₆X₇X₈X₉X₁₀X₁₁X₁₂X₁₃DX₁₄X₁₅X₁₆RX₁₇X₁₈X₂₇X₁₉X₂₀X₂₁X₂₂X₂₃X₂₄Y (SEQ ID NO:29) wherein X₁, X₈, X₁₃ and X₂₄ are each independently selected from the group consisting of L, I and V; X₂, X₄, X₅, X₆, X₇, X₉, X₁₀, X₁₁, X₁₂, X₁₄, X₁₅, X₁₆, X₁₇, X₁₈, X₁₉, X₂₀, X₂₁, X₂₂, and X₂₃ are each independently selected from the group consisting of any amino acid; X₃, X₂₅ and X₂₆ are each independently selected from the group consisting of any

35

-56-

- o amino acid and of no amino acid; and X_{27} is selected from the group consisting of L and Y.

5 15. A method of inducing serum antibodies which detect the toxins SPEA, SEA, SEB, and SED, comprising administering to a mammal, in a physiologically acceptable carrier, an immunologically sufficient amount of a peptide of claim 4.

16. The method of claim 15 wherein the peptide comprises the amino acid sequence
CMYGGVTEHEGNKKNVTQELDYKIRKYLVDNKKLYGC (SEQ ID NO: 8).

10 17. A method of inducing serum antibodies which inhibit blastogenesis of human mononuclear cells in the presence of any one of the toxins SEA, SEB, SEC, SEE, SPEA or SPEC comprising administering to a mammal, in a physiologically acceptable carrier, an immunologically sufficient amount of an antibody from a mammal immunized
15 with a peptide comprising a consensus amino acid sequence selected from the group consisting of $X_{25}X_{26}YGGX_1TX_2X_3X_4X_5N$ (SEQ ID NO:28) and
 $KX_6X_7X_8X_9X_{10}X_{11}X_{12}X_{13}DX_{14}X_{15}X_{16}RX_{17}X_{18}X_{27}X_{19}X_{20}X_{21}X_{22}X_{23}X_{24}Y$ (SEQ ID NO:29) wherein X_1 , X_8 , X_{13} and X_{24} are each independently
20 selected from the group consisting of L, I and V; X_2 , X_4 , X_5 , X_6 , X_7 , X_9 , X_{10} , X_{11} , X_{12} , X_{14} , X_{15} , X_{16} , X_{17} , X_{18} , X_{19} , X_{20} , X_{21} , X_{22} , and X_{23} are each independently selected from the group consisting of any amino acid; X_3 , X_{25} and X_{26} are each independently selected from the group consisting of any
25 amino acid and of no amino acid; and X_{27} is selected from the group consisting of L and Y.

18. A method of inducing serum antibodies which inhibit blastogenesis of human mononuclear cells in the presence of any one of the toxins SEA, SEB, SEC, SEE, SPEA
30 or SPEC comprising administering to a mammal, in a physiologically acceptable carrier, an immunologically sufficient amount of an antibody from a mammal immunized with a peptide of claim 4.

35 19. The method of claim 18 wherein the peptide

-57-

comprises the amino acid sequence

CMYGGVTEHEGNKKNVTVQELDYKIRKYLVDNKKLYGC (SEQ ID NO: 8).

20. A method of passively immunizing a mammal against the toxic effects of staphylococcal and streptococcal toxins comprising: administering in vivo an immunologically sufficient amount of an antibody containing composition wherein said antibody is derived from the immunization of antibody producing cells with a peptide comprising a consensus amino acid sequence selected from the group consisting of $X_{25}X_{26}YGGX_1TX_2X_3X_4X_5N$ (SEQ ID NO:28) and

$KX_6X_7X_8X_9X_{10}X_{11}X_{12}X_{13}DX_{14}X_{15}X_{16}RX_{17}X_{18}X_{27}X_{19}X_{20}X_{21}X_{22}X_{23}X_{24}Y$ (SEQ ID NO:29) wherein X_1 , X_8 , X_{13} and X_{24} are each independently selected from the group consisting of L, I and V; X_2 , X_4 , X_5 , X_6 , X_7 , X_9 , X_{10} , X_{11} , X_{12} , X_{14} , X_{15} , X_{16} , X_{17} , X_{18} , X_{19} , X_{20} , X_{21} , X_{22} , and X_{23} are each independently selected from the group consisting of any amino acid; X_3 , X_{25} and X_{26} are each independently selected from the group consisting of any amino acid and of no amino acid; and X_{27} is selected from the group consisting of L and Y.

21. A method of passively immunizing a mammal against the toxic effects of staphylococcal and streptococcal toxins comprising: administering in vivo an immunologically sufficient amount of an antibody containing composition wherein said antibody is derived from the immunization of antibody producing cells with a peptide of claim 4.

22. The method of claim 21 wherein the peptide comprises the amino acid sequence
CMYGGVTEHEGNKKNVTVQELDYKIRKYLVDNKKLYGC (SEQ ID NO: 8).

23. The method of claim 20 or claim 21 wherein the antibody composition is administered at a dose in the range of from about 1 mg/kg to about 10 mg/kg body weight of the mammal.

24. The method of any one of claims 9, 11, 14, 15, 17, 18, 20 and 21 wherein the mammal is a human.

-58-

25. A nucleic acid encoding a peptide comprising a consensus amino acid sequence selected from the group consisting of $X_{25}X_{26}YGGX_1TX_2X_3X_4X_5N$ (SEQ ID NO:28) and $KX_6X_7X_8X_9X_{10}X_{11}X_{12}X_{13}DX_{14}X_{15}X_{16}RX_{17}X_{18}X_{27}X_{19}X_{20}X_{21}X_{22}X_{23}X_{24}Y$ (SEQ ID NO:29) wherein X_1 , X_8 , X_{13} and X_{24} are each independently selected from the group consisting of L, I and V; X_2 , X_4 , X_5 , X_6 , X_7 , X_9 , X_{10} , X_{11} , X_{12} , X_{14} , X_{15} , X_{16} , X_{17} , X_{18} , X_{19} , X_{20} , X_{21} , X_{22} , and X_{23} are each independently selected from the group consisting of any amino acid; X_3 , X_{25} and X_{26} are each independently selected from the group consisting of any amino acid and of no amino acid; and X_{27} is selected from the group consisting of L and Y.

26. A nucleic acid encoding at least one amino acid sequence of claim 4.

27. A nucleic acid of claim 26 wherein the amino acid sequence encoded by said nucleic acid comprises CMYGGVTEHEGNKKNVTVQELDYKIRKYLVDNKKLYGC (SEQ ID NO: 8).

28. A host cell containing the nucleic acid of claim 25.

29. A host cell containing the nucleic acid of claim 26.

30. A method of inducing serum antibodies that bind staphylococcal enterotoxin and streptococcal exotoxin comprising administering to a mammal, in a physiologically acceptable carrier, a nucleic acid of claim 25 which produces an immunologically sufficient amount of the encoded peptide to elicit said antibodies.

31. A method of inducing serum antibodies that bind staphylococcal enterotoxin and streptococcal exotoxin comprising administering to a mammal, in a physiologically acceptable carrier, a nucleic acid of claim 26 which produces an immunologically sufficient amount of the encoded peptide to elicit said antibodies.

32. An antibody made by the method of any one of claims 9, 11, 14, 15, 17, and 19.

33. A method for detecting the presence of

-59-

- o staphylococcal or streptococcal toxin in a sample comprising contacting said sample with an antibody of claim 32 and detecting the antibody bound to said toxin.

34. A method for detecting the presence of antibodies to staphylococcal or streptococcal toxins in a sample comprising contacting said sample with a peptide
5 comprising a consensus amino acid sequence selected from the group consisting of $X_{25}X_{26}YGGX_1TX_2X_3X_4X_5N$ (SEQ ID NO:28) and $KX_6X_7X_8X_9X_{10}X_{11}X_{12}X_{13}DX_{14}X_{15}X_{16}RX_{17}X_{18}X_{27}X_{19}X_{20}X_{21}X_{22}X_{23}X_{24}Y$ (SEQ ID NO:29) wherein X_1 , X_8 , X_{13} and X_{24} are each independently selected from the group consisting of L, I and V; X_2 , X_4 ,
10 X_5 , X_6 , X_7 , X_9 , X_{10} , X_{11} , X_{12} , X_{14} , X_{15} , X_{16} , X_{17} , X_{18} , X_{19} , X_{20} , X_{21} , X_{22} , and X_{23} are each independently selected from the group consisting of any amino acid; X_3 , X_{25} and X_{26} are each independently selected from the group consisting of any amino acid and of no amino acid; and X_{27} is selected from
15 the group consisting of L and Y; and detecting the peptide bound to said antibodies.

35. A method for detecting the presence of antibodies to staphylococcal or streptococcal toxins in a sample comprising contacting said sample with a peptide of
20 claim 4 and detecting the peptide bound to said antibodies.

36. A kit for detecting the presence of staphylococcal or streptococcal toxins in a sample comprising an antibody of claim 32.

25 37. A kit for detecting the presence of antibodies to staphylococcal or streptococcal toxins in a sample comprising a peptide comprising a consensus amino acid sequence selected from the group consisting of $X_{25}X_{26}YGGX_1TX_2X_3X_4X_5N$ (SEQ ID NO:28) and
30 $KX_6X_7X_8X_9X_{10}X_{11}X_{12}X_{13}DX_{14}X_{15}X_{16}RX_{17}X_{18}X_{27}X_{19}X_{20}X_{21}X_{22}X_{23}X_{24}Y$ (SEQ ID NO:29) wherein X_1 , X_8 , X_{13} and X_{24} are each independently selected from the group consisting of L, I and V; X_2 , X_4 , X_5 , X_6 , X_7 , X_9 , X_{10} , X_{11} , X_{12} , X_{14} , X_{15} , X_{16} , X_{17} , X_{18} , X_{19} , X_{20} , X_{21} , X_{22} , and X_{23} are each independently selected from the
35

-60-

- o group consisting of any amino acid; X_3 , X_{25} and X_{26} are each independently selected from the group consisting of any amino acid and of no amino acid; and X_{27} is selected from the group consisting of L and Y.

5 38. A kit for detecting the presence of antibodies to staphylococcal or streptococcal toxins in a sample comprising a peptide of claim 4.

10

15

20

25

30

35

1/4


Figure 1.

TOXIN	REGION 1		REGION 2	
PEPs:	<u>CMYGGVTEHEGN</u>		<u>KKNVTVQELDYKIRKYLVDNKKLY</u>	
SEA	130	CMYGGVTLHDNN 141	171	KKNVTVQELDLQARRYLQEKYNLY 194
SEB	140	CMYGGVTEHNGN 151	179	KKKVTAQELDYLTRHYLVKNKKLY 202
SEC	137	CMYGGITKHEGN 148	178	KKSVTAQELDIKARNFLINKKNLY 201
SED	131	CTYGGVTPHEGN 142	172	KKNVTVQELDAQARRYLQKDLKLY 195
SEE	130	CMYGGVTLHDNN 141	171	KKEVTVQELDLQARHYLHGKFGLY 194
SEH	116	CLYGGITL.NSE 126	151	KKNVTLQELDIKIRKILSDKYKIY 174
SPEA	128	CIYGGVTNHEGN 139	167	KKMVTQELDYKVRKYLTDNKQLY 190
SPEC	112	YIYGGITPAQNN 123	151	KDIVTFQEIDFKIRKLYMDNYKIY 174
SSA	134	CMYGGVTEHHRN 145	174	KKQVTVQELDCCKTRKILVSRKNLY 197
TSST1			161	KKQLAISTLDFEIRHQLTQIHGLY 184

THIS PAGE BLANK (USPTO)


Figure 2:

ELISA Titers of Rabbit Anti-peptide Antibodies
Against Peptide Polymer



THIS PAGE BLANK (USPTO)


Figure 3.



THIS PAGE BLANK (USPTO)

Figure 4

Inhibition of Blastogenesis of Human Mononuclear Cell Population
Using Anti-peptide Antibody and Various Toxins



THIS PAGE BLANK (USPTO)

INTERNATIONAL SEARCH REPORT

In. .national Application No

PCT/US 98/06663

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07K14/31 C07K16/12 C12N15/31 G01N33/577 A61K39/085
A61K39/09

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 36366 A (NATIONAL JEWISH CENTER FOR IMMUNOLOGY) 21 November 1996 see the whole document and specially pages 95-96 ---	1,4,5,8, 11,15, 21,26, 29,31, 35,38
X	WO 91 10680 A (D.S.TERMAN AND J.L. STONE) 25 July 1991 see page 15-16; tables 2,6 see figures 1,2 --- -/--	1,4,5,8, 11,15, 21,26, 29,31, 35,38

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

5 August 1998

Date of mailing of the international search report

21/08/1998

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Mateo Rosell, A.M.

INTERNATIONAL SEARCH REPORT

In. .ational Application No

PCT/US 98/06663

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 93 24136 A (D.S.TERMAN AND J.L.STONE) 9 December 1993</p> <p>see abstract see page 5, line 26 - page 6, line 14; tables 2,6 see page 14-17</p>	<p>1,4,5,8, 11,15, 21,26, 29,31, 35,38</p>
X	<p>WO 94 20124 A (D.S.TERMAN) 15 September 1994</p> <p>see page 7, line 17 - page 8, line 10; tables 2,6</p>	<p>1,4</p>
A	<p>WO 96 40930 A (UNIV MINNESOTA) 19 December 1996</p> <p>see the whole document and specially sequence disclosed in pages 77-79</p>	<p>1,4,5,8, 11,15, 21,26, 29,31, 35,38</p>
A	<p>WO 94 25483 A (UNIV BRITISH COLUMBIA) 10 November 1994</p> <p>cited in the application See the whole document and specially SEQ.ID.N.4</p>	<p>1,17-19</p>
A	<p>VAN DER BUSSCHE ET AL.,: "Molecular evolution of the staphylococcal and streptococcal pyrogenic toxin gene family" MOLECULAR PHYLOGENETICS AND EVOLUTION, vol. 2, no. 4, 1993, pages 281-292, XP002073580</p> <p>cited in the application see the whole document and specially figures 1 and 2</p>	<p>1</p>

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/06663

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 9-24 AND 30-31
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim(s) 9-24 AND 30-31
is(are) directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/06663

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9636366 A	21-11-1996	US 5705151 A AU 5801696 A EP 0850071 A	06-01-1998 29-11-1996 01-07-1998
WO 9110680 A	25-07-1991	CA 2078003 A EP 0511306 A US 5728388 A	18-07-1991 04-11-1992 17-03-1998
WO 9324136 A	09-12-1993	AU 4526593 A WO 9420124 A US 5728388 A	30-12-1993 15-09-1994 17-03-1998
WO 9420124 A	15-09-1994	AU 6358994 A CA 2157379 A EP 0705104 A US 5728388 A AU 4526593 A WO 9324136 A	26-09-1994 15-09-1994 10-04-1996 17-03-1998 30-12-1993 09-12-1993
WO 9640930 A	19-12-1996	AU 6278296 A EP 0832241 A	30-12-1996 01-04-1998
WO 9425483 A	10-11-1994	AU 6760294 A	21-11-1994